

Double Higgs production in the Standard Model with extended scalar sector

1408.0184, 1503.01618

S. I. Godunov, M. I. Vysotsky, A. N. Rozanov,
E. V. Zhemchugov

International conference on
particle physics and astrophysics
October 5–10, Moscow, Russia

Motivation

- ▶ A scalar with mass 125 GeV has been discovered in 2012.
- ▶ In order to confirm that this is the Standard Model Higgs boson, its couplings have to be measured.
- ▶ Triple coupling: $g_{hhh} \sim \frac{m_h^2}{v}$. It can be measured in the $pp \rightarrow hh$ process.
- ▶ The Standard Model prediction for the $pp \rightarrow hh$ cross section is 40 fb for $\sqrt{s} = 14$ TeV. That can only be measured at HL-LHC.
- ▶ What if there are other scalar particles?

Isosinglet

arXiv:1503.01618

Scalar sector:

$$\Phi = \begin{pmatrix} \phi^+ \\ \frac{1}{\sqrt{2}}(v_\Phi + \phi + i\eta) \end{pmatrix}, \quad X = v_X + \chi$$

Potential:

$$V_1(\Phi, X) = -\frac{1}{2}m_\Phi^2\Phi^\dagger\Phi + \frac{\lambda}{2}(\Phi^\dagger\Phi)^2 + \frac{1}{2}m_X^2X^2 + \mu\Phi^\dagger\Phi X$$


Mixing:

$$\begin{pmatrix} h \\ H \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi \\ \chi \end{pmatrix}.$$

1. $\left. \frac{\partial V_1}{\partial \phi} \right|_{\phi=0, \chi=0} = 0,$
2. $\left. \frac{\partial V_1}{\partial \chi} \right|_{\phi=0, \chi=0} = 0,$
3. $v_\Phi = 246$ GeV from the Fermi coupling in muon decay.
4. h is associated with the SM-like higgs, so $m_h = 125$ GeV.

Remaining model parameters: $\sin \alpha$ and m_H .


Bounds from electroweak precision observables and μ measurements:





$$\mu_i = \frac{\sigma(pp \rightarrow h) \cdot \text{Br}(h \rightarrow f_i)}{(\sigma(pp \rightarrow h) \cdot \text{Br}(h \rightarrow f_i))_{\text{SM}}} = \cos^2 \alpha$$


ATLAS: $\mu = 1.30^{+0.18}_{-0.17}$, CMS: $\mu = 1.00^{+0.14}_{-0.13}$.

$$R \equiv \frac{\sigma(pp \rightarrow H) \text{Br}(H \rightarrow ZZ)}{(\sigma(pp \rightarrow h) \text{Br}(h \rightarrow ZZ))_{\text{SM}}} = \frac{\sin^4 \alpha}{\sin^2 \alpha + \frac{\Gamma(H \rightarrow hh)}{\Gamma_{\text{SM}}}}$$





CMS PAS HIG-13-031



$$\sigma(pp \rightarrow H \rightarrow hh) = \sigma(pp \rightarrow h)_{\text{SM}} \cdot \sin^2 \alpha \cdot \mathcal{B}(H \rightarrow hh)$$

Isotriplet

J. Exp. Theor. Phys. 120 (2015) 3, 369–375
arXiv:1408.0184

Scalar sector:

$$\Phi = \begin{bmatrix} \phi^+ \\ \frac{1}{\sqrt{2}}(v_\Phi + \phi + i\eta) \end{bmatrix}, \quad \Delta = \frac{\vec{\Delta}\vec{\sigma}}{\sqrt{2}} = \begin{bmatrix} \delta^+/\sqrt{2} & \delta^{++} \\ \frac{1}{\sqrt{2}}(v_\Delta + \delta + i\rho) & -\delta^+/\sqrt{2} \end{bmatrix}$$

Potential:

$$V(\Phi, \Delta) = -\frac{1}{2}m_\Phi^2(\Phi^\dagger\Phi) + \frac{\lambda}{2}(\Phi^\dagger\Phi)^2 \\ + m_\Delta^2 \text{tr}[\Delta^\dagger\Delta] + \frac{\mu}{\sqrt{2}}(\Phi^T i\sigma^2 \Delta^\dagger\Phi + \text{h.c.})$$

Mixing:

$$\begin{pmatrix} h \\ H \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi \\ \delta \end{pmatrix}.$$

$$m_h = 125 \text{ GeV}.$$

Custodial symmetry breaking:

$$\left. \begin{aligned} m_W^2 &= \frac{g^2}{4}(v_\Phi^2 + 2v_\Delta^2) \\ m_Z^2 &= \frac{\bar{g}^2}{4}(v_\Phi^2 + 4v_\Delta^2) \end{aligned} \right\} \Rightarrow \frac{m_W}{m_Z} \approx \left(\frac{m_W}{m_Z} \right)_{\text{SM}} \left(1 - \frac{v_\Delta^2}{v_\Phi^2} \right)$$

$$\left(\frac{m_W}{m_Z \cos \theta_W} \right)_{\text{SM}} = 1.00040 \pm 0.00024 \Rightarrow v_\Delta < 5 \text{ GeV (at } 3\sigma \text{ level)}$$

In the following we assume $v_\Delta = 5 \text{ GeV}$.


$$v_\Phi^2 + 2v_\Delta^2 = \frac{1}{\sqrt{2}G_F} = (246 \text{ GeV})^2 \Rightarrow v_\Phi \approx 246 \text{ GeV}.$$

With $v_\Delta \ll v_\Phi$ we get $\sin \alpha \approx \frac{2v_\Delta}{v_\Phi} \ll 1$.

Remaining model parameters: m_H . We will consider the case of $m_H = 300 \text{ GeV}$.

▶ $H \rightarrow WW$ is suppressed as $(m_h/m_H)^4 \approx 0.03$.

▶ $\mathcal{B}(H \rightarrow hh) \approx 0.8$.



Higgs boson double production cross section at the LHC for $\sqrt{s} = 14$ TeV

Mass, GeV	the SM h		H
	125	300	300
$\sigma(gg \rightarrow h, H),^1$ pb	50(5)	11(1)	$25(2) \cdot 10^{-3}$
$\sigma(gg \rightarrow t\bar{t} + h, H),^1$ pb	0.61(6)	0.051(5)	$12(1) \cdot 10^{-6}$
$\sigma(W^+W^- \rightarrow h, H),^2$ pb	3.272(4)	1.053(1)	$76.8(1) \cdot 10^{-6}$
$\sigma(ZZ \rightarrow h, H),^2$ pb	1.087(1)	0.365(1)	$365(1) \cdot 10^{-6}$
$\sigma(W^* \rightarrow Wh, WH),^2$ pb	0.150(6)	0.068(3)	$5.0(2) \cdot 10^{-6}$
$\sigma(Z^* \rightarrow Zh, ZH),^2$ pb	0.88(5)	0.042(2)	$42(2) \cdot 10^{-6}$

$$\begin{aligned} \sigma(pp \rightarrow hh) &= 40 \text{ fb (SM)} + 25 \text{ fb (H production)} \cdot 0.8 \text{ (branching)} \\ &= 60 \text{ fb} \end{aligned}$$

¹The SM values are from *Handbook of LHC Higgs cross sections*, CERN-2011-002.

²The SM values were obtained with the help of the program HAWK at LO + QCD without electroweak corrections.

The Georgi-Machacek model

$$\Phi = \begin{bmatrix} \phi^{0*} & \phi^+ \\ -\phi^- & \phi^0 \end{bmatrix}, \quad \Delta = \begin{bmatrix} \delta^{0*} & \xi^+ & \delta^{++} \\ -\delta^- & \xi^0 & \delta^+ \\ \delta^{--} & -\xi^- & \delta^0 \end{bmatrix}.$$

$$\phi^0 = \frac{1}{\sqrt{2}}(v_\Phi + \phi + i\chi), \quad \delta^0 = \frac{1}{\sqrt{2}}(v_\Delta + \delta + i\eta), \quad \xi^0 = v_\xi + \xi.$$

$$m_W^2 = \frac{g^2}{4}(v_\Phi^2 + 2v_\Delta^2 + 2v_\xi^2), \quad m_Z^2 = \frac{\bar{g}^2}{4}(v_\Phi^2 + 4v_\Delta^2).$$

$$(v_\Phi^2 + 2v_\Delta^2 + 2v_\xi^2) = \frac{1}{\sqrt{2}G_F} = (246 \text{ GeV})^2.$$



When $v_\xi = v_\Delta$, the bound on v_Δ from the gauge bosons mass ratio is relieved. LHC measurements allow v_Δ up to 50 GeV. In this case for $m_H = 300 \text{ GeV}$ and $\sqrt{s} = 14 \text{ TeV}$ $\sigma(gg \rightarrow H) = 1.4 \text{ pb}$ with $\text{Br}(H \rightarrow hh) = 98\%$ and $\text{Br}(H \rightarrow ZZ) = 0.6\%$.

Conclusions

- ▶ **Isosinglet:** $\sigma(pp \rightarrow hh)$ can be over 0.4 pb for $\sqrt{s} = 14$ TeV.
 - ▶ $H \rightarrow ZZ$ is the golden mode.
- ▶ **Isotriplet:** $\sigma(pp \rightarrow hh) = 60$ fb for $\sqrt{s} = 14$ TeV and $m_H = 300$ GeV.
 - ▶ $H \rightarrow ZZ$ is the golden mode; $H \rightarrow W^+ W^-$ is heavily suppressed.
 - ▶ Custodial symmetry is broken.
- ▶ **The Georgi-Machacek model:** $\sigma(pp \rightarrow hh)$ can be as high as 1.4 pb for $\sqrt{s} = 14$ TeV.
 - ▶ When $m_H \approx 300$ GeV $\mathcal{B}(H \rightarrow hh) = 0.98$.
 - ▶ Custodial symmetry is preserved.



Thank you for your attention

$$R \equiv \frac{\sigma(pp \rightarrow H) \text{Br}(H \rightarrow ZZ)}{(\sigma(pp \rightarrow h) \text{Br}(h \rightarrow ZZ))_{\text{SM}}} = \frac{\sin^4 \alpha}{\sin^2 \alpha + \frac{\Gamma(H \rightarrow hh)}{\Gamma_{\text{SM}}}}$$



CERN-PH-EP-2015-154

$$R \equiv \frac{\sigma(pp \rightarrow H) \text{Br}(H \rightarrow ZZ)}{(\sigma(pp \rightarrow h) \text{Br}(h \rightarrow ZZ))_{\text{SM}}} = \frac{\sin^4 \alpha}{\sin^2 \alpha + \frac{\Gamma(H \rightarrow hh)}{\Gamma_{\text{SM}}}}$$



CMS PAS HIG-13-002.