Contribution ID : 97

Visible, invisible and trapped ghosts in wormholes and black universes

Friday, 9 October 2015 13:15 (15)

S.V. Bolokhov, K.A. Bronnikov, P.A. Korolyov, M.V. Skvortsova

We construct explicit examples of globally regular static, spherically symmetric solutions in general relativity with scalar and electromagnetic fields, describing (1) traversable wormholes with flat and AdS asymptotics and (2) regular black holes, in particular, black universes. (A black universe is a regular black hole with an expanding, asymptotically isotropic space-time beyond the horizon.) The existence of such objects requires invoking scalars with negative kinetic energy (phantoms", orghosts"), which are not observed under usual physical conditions. To account for that, the so-called trapped ghosts" were previously introduced, i.e., scalars whose kinetic energy is only negative in a restricted strong-field region of space-time and positive outside it. This approach leads to certain problems, including instability (as we illustrate by derivation of an effective potential for spherical pertubations of such systems). To avoid that, we use for model construction what we callinvisible ghosts", i.e., phantom scalar fields sufficiently rapidly decaying in the weak-field region. The resulting configurations contain different numbers of Killing horizons, from zero to four. Their global structure has been studied with the aid of Carter-Penrose diagrams.

Presentation type

Section talk (10+5 min)

Primary author(s): Prof. BRONNIKOV, Kirill (VNIIMS)

Co-author(s) : Dr. SKVORTSOVA, Milena (VNIIMS, Moscow); Mr. KOROLYOV, Pavel (PFUR, Moscow); Dr. BOLOKHOV, Sergei (PFUR, Moscow)

Presenter(s) : Mr. KOROLYOV, Pavel (PFUR, Moscow)

Session Classification : Nuclear physics and particle physics - parallel XI

Track Classification : Nuclear physics and particle physics