

# Yields of p, d, t formed in stopped pion absorption by intranuclear clusters

*Tuesday, 6 October 2015 13:45 (15)*

Results on the measurement of the spectra of p, d, t formed in the reaction of stopped pion absorption by atomic nuclei from the wide mass number range ( $6 < A < 209$ ) are presented. Phenomenological model developed by us [1, 2] had allowed to calculate contributions of cluster absorption, preequilibrium and evaporation processes in the particle formation. A-dependences of primary particles formed in the cluster absorption process are obtained. Analytical expressions are developed that allow to describe A-dependences successfully in the investigated mass number range.

1. L.Yu.Korotkova et al. // Bull. of RAS.: Phys. 2012. V. 76. P. 446.
2. Yu.B.Gurov et al. // Bull. of RAS.: Phys. 2013. V. 77. P. 415.

## Presentation type

Section talk (10+5 min)

**Primary author(s) :** Mr. PRITULA, Roman (MEPhI)

**Co-author(s) :** Dr. CHERNYSHEV, Boris (National Research Nuclear University MEPhI); Mrs. KOROTKOVA, Larisa (National Research Nuclear University MEPhI); Dr. LAPUSHKIN, Sergei (National Research Nuclear University "MEPhI"); Mrs. LEONOVA, Tatiana (National Research Nuclear University "MEPhI"); Dr. SCHURENKOVA, Tatiana (National Research Nuclear University "MEPhI"); Dr. GUROV, Yurii (National Research Nuclear University "MEPhI")

**Presenter(s) :** Mr. PRITULA, Roman (MEPhI)

**Session Classification :** Nuclear physics and particle physics - parallel I

**Track Classification :** Nuclear physics and particle physics