

# Kinks in the relativistic model with logarithmic nonlinearity

*Monday, 22 October 2018 15:40 (150)*

We study the properties of a certain class of solutions of a relativistic model with the logarithmic nonlinearity. We show that such a model have two types of solutions: topologically trivial (gaussons) and topologically non-trivial (kinks). For the kink-antikink scattering, we have found a critical value of the initial velocity  $v_{cr}$ , which separates two different scenarios of scattering. At the initial velocities  $v_{in} < v_{cr}$ , the kinks form a bound state, which then decays slowly. At  $v_{in} > v_{cr}$ , the kinks collide, bounce and eventually escape to infinities. During this process, the higher initial velocity is, the greater is the elasticity of the collision. We also study excitation spectrum of the kink solution.

**Primary author(s) :** Dr. GANI, Vakhid (National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), 115409 Moscow, Russia); Dr. ZLOSHCHASTIEV, Konstantin (Institute of Systems Science, Durban University of Technology, P.O. Box 1334, Durban 4000, South Africa)

**Presenter(s) :** Dr. GANI, Vakhid (National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), 115409 Moscow, Russia)

**Session Classification :** Poster session and coffee-buffet

**Track Classification :** Particle physics: hep theory