



Contribution ID : 263

Type : Oral talk

## Isoscalar giant monopole resonance in the Ca isotopes

Wednesday, 30 November 2022 19:45 (15)

A comprehensive analysis of the isoscalar giant monopole resonance (ISGMR) has long been a subject of extensive theoretical and experimental research [1,2]. The ISGMR properties are presently an important problem not only from the nuclear structure point of view [2,3] but also because of the special role they play in many astrophysical processes such as prompt supernova explosions [4] and the interiors of neutron stars [5].

The random phase approximation (RPA) with the Skyrme-type energy density functional (EDF) is the most widely used theoretical model for describing the ISGMR [2,3]. The study of the monopole strength distribution in the region of giant resonance involves taking into account a coupling between the simple particle-hole excitations and more complicated (two- and three-phonons) configurations [3,6]. The main difficulty is that the complexity of calculations beyond standard RPA increases rapidly with the size of the configuration space, and one has to work within limited spaces. Using a finite rank separable approximation for the residual particle-hole interaction derived from the Skyrme EDF one can overcome this numerical problem [7,8].

In the present report, we discuss the effects of the coupling between one-, two-, and three-phonon terms in the wave functions on the monopole strength distribution in the double-magic nuclei  $^{40,48}\text{Ca}$ . Using the same set of parameters, we describe available experimental data [9,10]. The effects of the phonon-phonon coupling (PPC) lead to a redistribution of the main monopole strength to lower energy states and into higher energy tail [11]. The PPC predictions of the fine structure of the ISGMR in the Ca isotopes are in good agreement with the fine structure which is extracted from experimental data analysis [12].

This work was supported by the Russian Science Foundation (Grant No. RSF-21-12-00061).

1. J.P. Blaizot, Phys. Rev. 64, 171 (1980).
2. U. Garg, and G. Colò, Prog. Part. Nucl. Phys. 101, 55 (2018).
3. N.N. Arsenyev, and A.P. Severyukhin, Universe. 7, 145 (2021).
4. H.A. Bethe, Rev. Mod. Phys. 62, 801 (1990).
5. N.K. Glendenning, Phys. Rev. Lett. 57, 1120 (1986).
6. V.G. Soloviev, Theory of Atomic Nuclei: Quasiparticles and Phonons. Bristol/Philadelphia 1992.
7. N.V. Giai, Ch. Stoyanov, and V.V. Voronov, Phys. Rev. C. 57, 1204 (1998).
8. A.P. Severyukhin, V.V. Voronov, and N.V. Giai, Eur. Phys. Jour. A. 22, 397 (2004).
9. K. Howard et al., Phys. Lett. B. 801, 135185 (2020).
10. S.D. Olorunfunmi et al., Phys. Rev. C. 105, 054319 (2022).
11. N.N. Arsenyev, and A.P. Severyukhin, Phys. At. Nucl. 85, (2022) in press.
12. S.D. Olorunfunmi et al., in preparation.

**Primary author(s) :** ARSENYEV, Nikolay (Bogoliubov Laboratory of Theoretical Physic, Joint Institute for Nuclear Research); Dr. SEVERYUKHIN, Alexey (Bogoliubov Laboratory of Theoretical Physic, Joint Institute for Nuclear Research)

**Presenter(s)** : ARSENYEV, Nikolay (Bogoliubov Laboratory of Theoretical Physic, Joint Institute for Nuclear Research)

**Session Classification** : Nuclear Physics

**Track Classification** : Nuclear physics