



Contribution ID : 290

Type : Oral talk

Correction to Bjorken energy density calculations for central A-A collisions

Thursday, 1 December 2022 19:30 (15)

O.M. Shaposhnikova¹, A.A. Marova², G.A. Feofilov^{2,*}

¹ St. Petersburg Governor's Physics and Mathematics Lyceum No. 30

² Federal State Budgetary Educational Institution of Higher Education "Saint-Petersburg State University", Saint Petersburg, 199034 Russia

*E-mail: g.feofilov@spbu.ru

Abstract:

We consider a correction to the Bjorken energy density [1] calculations [2-6] where the information is usually taken from the nucleus-nucleus collisions events selected in a certain centrality class. The last one is usually defined in terms of some observable (multiplicity of particles, number of participating nucleons, transverse energy) used as a proxy to the impact parameter b . However, due to the natural dominance of peripheral collisions in any class of centrality, the relevant mean values of the impact parameter will be always shifted towards some higher values. For example, it is possible to show that in case of 0-5% centrality class of A-A collisions, the value is considerably shifted from 0 (~ 2 fm), therefore in this case the corresponding value of ST - the transverse interaction area of two overlapping colliding disks, should not be taken to be that of a circle. It appears to be smaller, thus resulting in the higher values of energy density finally calculated for a given class.

We present our estimates of the Bjorken energy density in 0-5% centrality classes events in a broad range of heavy-ion collision energies (from SPS to the LHC) using the available experimental information on mean multiplicity of charged particles and mean transverse momenta. We make the analytical approximation of the updated Bjorken energy excitation function, obtained in our study for these very central A-A collisions, and we compare it to the previously obtained ones [7,8]. We present also and discuss the relevant energy dependences of pion, kaon and proton contributions to the Bjorken energy density that are also estimated in our study.

This work was supported by St. Petersburg State University (project № 93025435). References:

- 1) J. D. Bjorken, Phys. Rev. D 27, 140 (1983).
- 2) B. I. Abelev, M. M. Aggarwal et al. Phys.Rev. C 79, 034909 (2009).
- 3) K. Adcox, S. S. Adler et al. arXiv:nucl-ex/0104015 v2(2001).
- 4) B. Abelev et al., Phys.Rev. C 88, 044909 (2013).
- 5) D. Adhikari, H. Albatineh et al., Phys.Rev.Lett 126, 172502 (2021).
- 6) S. S. Adler, S. Afanasiev et al., Phys.Rev. C 71, 034908 (2005).
- 7) T. Mendenhall, Z.-W. Lin, Phys. Rev. C 103, 024907 (2021).
- 8) R.Sahoo et al., Advances in High Energy Physics, v. 2015, 612390 (2015).

Primary author(s): SHAPOSHNIKOVA, Olga

Co-author(s) : Dr. FEOFILOV, Grigory (Saint-Petersburg State University); MAROVA, Aleksandra (Saint Petersburg State University)

Presenter(s) : SHAPOSHNIKOVA, Olga

Session Classification : High Energy Physics: Theory

Track Classification : High energy physics: theory