

Investigation of the correlation between mean transverse momentum and anisotropic flow at NICA energy range

Idrisov Dim¹, Peter Parfenov^{1,2}

1. National Research Nuclear University (MEPhI), Moscow, Russia
2. Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia

The 6th International Conference on Particle Physics and Astrophysics

MEPhI, Moscow


29.11-02.12.2022

Outline


- Introduction
- Method for the transverse momentum-flow correlations measurements
- Comparison with published data
- The results at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV for different models
- Summary and outlook

Elliptic flow at NICA energies

$$\frac{dN}{d\varphi} \sim 1 + \sum_{n=1} v_n \cos[n(\varphi - \Psi_n)],$$




$$v_2\{2\} = \sqrt{\langle e^{i2(\varphi_1 - \varphi_2)} \rangle}, \quad v_2\{4\} = \sqrt{4 \langle e^{i2(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4)} \rangle - 2 \langle e^{i2(\varphi_1 - \varphi_2)} \rangle^2}$$



- v_2 is sensitive to the properties of strongly interacting matter:

- At $\sqrt{s_{NN}} \geq 7.7$ GeV pure string/hadronic cascade models underestimate v_2 – need hybrid models with QGP phase (vHLE+UrQMD, AMPT SM)



- Relative v_2 fluctuations ($v_2\{4\}/v_2\{2\}$) observed by STAR experiment can be reproduced in the string/cascade models (UrQMD, SMASH) and hybrid model (AMPT SM, vHLE+UrQMD)
- Dominant source of v_2 fluctuations: **participant eccentricity fluctuations** in the initial geometry

The correlation coefficient

The correlation coefficient defined as

$$\rho(v_2^2, [p_T]) = \frac{\text{cov}(v_2^2, [p_T])}{\sqrt{\text{var}(v_2^2)}\sqrt{c_k}}$$

where $\text{var}(v_2^2)_{\text{dyn}} = \langle v_2^4 \rangle - \langle v_2^2 \rangle^2 = \langle \langle 4 \rangle \rangle_{A,C} - \langle \langle 2 \rangle \rangle_{A,C}^2$

$$\langle \langle 2 \rangle \rangle_{A,C} = \left\langle \left\langle e^{i \cdot 2(\phi_1^A - \phi_2^C)} \right\rangle \right\rangle = \frac{Q_{2,A} Q_{2,C}^*}{M_A M_C},$$

$$\langle \langle 4 \rangle \rangle_{A,C} = \left\langle \left\langle e^{i \cdot 2(\phi_1^A + \phi_2^A - \phi_3^C - \phi_4^C)} \right\rangle \right\rangle = \frac{(Q_{2,A}^2 - Q_{4,A})(Q_{2,C}^2 - Q_{4,C})^*}{M_A(M_A - 1)M_C(M_C - 1)}$$

$$Q_{n,A/C} = \sum_k e^{i \cdot n \phi_k^{A/C}} - \text{flow vector for A/C sub event}$$

$M_{A/C}$ - multiplicity of particles

$$-1 < \eta < -0.35 \quad |\eta| < 0.35 \quad 0.35 < \eta < 1$$



to suppress non-flow effects,
the two sub-events method was used

In the study were used charged particles
with $0.2 < p_T < 2.0 \text{ GeV/c}$

The correlation coefficient 2

The variance of the mean transvers momentum, taking into account autocorrelations, is defined as

$$c_k = \left\langle \frac{1}{M_B(M_B - 1)} \sum_B \sum_{B' \neq B} (p_{T,B} - \langle [p_T] \rangle) (p_{T,B'} - \langle [p_T] \rangle) \right\rangle$$


where $[p_T] = \frac{\sum_{i=1}^{M_B} p_{T,i}}{M_B}$

to suppress non-flow and autocorrelation effects

in the $\text{cov}(v_2^2, [p_T])$ the three-subevents method was used

$$\text{cov}(v_2^2, [p_T]) = \left\langle \frac{\sum_{A,C} e^{i \cdot 2(\phi_1^A - \phi_2^C)} \sum_B (p_{T,B} - \langle [p_T] \rangle)}{M_A M_C M_B} \right\rangle$$



$$-1 < \eta < -0.35 \quad |\eta| < 0.35 \quad 0.35 < \eta < 1$$



Motivation of the work

- The $\rho(v_2^2, [p_T])$ is sensitive to initial state and its entropy density profile
- The $\text{cov}(v_2^2, [p_T])$ and $\text{var}(v_2^2)$ are sensitive to η/s
- The precise set of measurements for $\text{var}([p_T])$, $\text{var}(v_2^2)$, $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ as a function of beam-energy and centrality, could aid precision extraction of the temperature and baryon chemical-potential dependence of η/s



Centrality for Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV in vHLLE+UrQMD



The reasonable fit quality and good agreement of the impact parameter distribution with the model data. For centrality determination the Inverse Bayes approach was used.



Comparison of correlation coefficient with published results

The published data taken from: Niseem Magdy et. al. Published in: Phys.Rev.C 105 (2022) 4, 044901




Filled red squares: multiplicity-based centrality
Open black squares: b-based centrality

- A good agreement between published data and results with b-based centrality
- The $\text{cov}(v_2^2, [p_T])$ is sensitive to the multiplicity fluctuations







The $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ depend on the centrality determination method.

Elliptic flow and its fluctuations at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV







The flow fluctuations are model independent and decrease with decreasing energy.

The transverse momentum-flow correlations at $\sqrt{s_{NN}}=11.5$ GeV



- $\rho(v_2^2, [p_T])$ decreases in the most central collisions due to the eccentricity decreases faster compared to changes in elliptic area.
- $\rho(v_2^2, [p_T])$ from vHLE+UrQMD and UrQMD are consistent with each other due to the same initial state
 - $\rho(v_2^2, [p_T])$ is sensitive to initial state
- $\text{cov}(v_2^2, [p_T])$ from vHLE+UrQMD and AMPT are consistent due to QGP phase
 - $\text{cov}(v_2^2, [p_T])$ is sensitive to thermalization (η/s , etc.)

The transverse momentum-flow correlations at $\sqrt{S_{NN}}=7.7$ GeV







- The same trends as for $\sqrt{s_{NN}}=11.5$ GeV
- The $\text{var}(v_2^2)$ decrease with decreasing energy
- More statistics are needed to get more accurate results

Summary and outlook

- A good agreement between published data and results for vHLE+UrQMD at $\sqrt{s_{NN}} = 62.4$ GeV with b-based centrality for $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$
 - The $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ depend on the centrality determination method
- The results at $\sqrt{s_{NN}} = 7.7$ and 11.5 GeV for AMPT, UrQMD, and vHLE+UrQMD
 - $\rho(v_2^2, [p_T])$ from vHLE+UrQMD and UrQMD are consistent with each other due to the same initial state
 - $\text{cov}(v_2^2, [p_T])$ from vHLE+UrQMD and AMPT are consistent due to QGP phase simulation
 - $\rho(v_2^2, [p_T])$ decreases in the most central collisions
 - for the first time, results were obtained at $\sqrt{s_{NN}} = 7.7$ and 11.5 GeV
- Investigate beam-energy and event-shape dependence of the $v_2^2 - [p_T]$ correlation using vHLE+UrQMD model
- Study sensitivity of $v_2^2 - [p_T]$ correlation to different equation of states in models within mean-field approach at lower beam energies





Thank you for your attention!

The results for vHLE+UrQMD at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV







- The $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ changes weakly at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV
- The $\sqrt{c_k}/\langle p_T \rangle$ increases with decreasing energy.

The results for UrQMD at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV



- Do the $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ increases with decreasing energy due to non-flow effects?

The results for AMPT at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV





- The set of measurements for $\text{var}([p_T])$, $\text{var}(v_2^2)$, $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ changes weakly at $\sqrt{S_{NN}}=7.7$ and 11.5 GeV

Summary and outlook

- A good agreement between published data and results for vHLE+UrQMD at $\sqrt{S_{NN}} = 62.4$ GeV with b-based centrality for $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$
 - The $\text{cov}(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ depend on the centrality determination method
- The results at $\sqrt{S_{NN}} = 7.7$ and 11.5 GeV for AMPT, UrQMD, and vHLE+UrQMD
 - The $\rho(v_2^2, [p_T])$ vs. centrality for vHLE+UrQMD at $\sqrt{S_{NN}} = 7.7$ and 11.5 shows the similar trends as for BES energies.
 - $\rho(v_2^2, [p_T]) < 0$ for the most central collisions in UrQMD and AMPT models at $\sqrt{S_{NN}} = 7.7$ and 11.5 GeV
- Investigate beam-energy and event-shape dependence of the $v_3^2 - [p_T]$ correlation using vHLE+UrQMD model
- Study sensitivity of $v_2^2 - [p_T]$ correlation to different equation of states in models within mean-field approach at lower beam energies



Transverse momentum-flow correlations

Models	Initial stage conditions	η/s
AMPT	Glauber-like & SM-ON	0.10
AMPT	Glauber-like & SM-OFF	0.10
EPOS	Described in terms of flux tubes computed based on Gribov-Regge multiple scattering theory	0.08
Hydro	Woods-Saxon distributions Glauber-like Initial conditions	(BES)




- The $cov(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$ show agreement between AMPT (SM) and EPOS
- Smaller $cov(v_2^2, [p_T])$ and $\rho(v_2^2, [p_T])$, from AMPT without SM

Transverse momentum-flow correlations



- The $\text{cov}(v_2^2, [p_T])$ decreases with η/s
- The $\rho(v_2^2, [p_T])$, show weak dependance on η/s




The transverse momentum-flow correlations dependence on beam energy in vHLE+UrQMD model



- The $\text{cov}(v_2^2, [p_T])$ decreases with beam energy
- The $\rho(v_2^2, [p_T])$, show weak dependance on beam energy

Comparison of elliptic flow measurements with published results




Filled red squares: multiplicity-based centrality
Open black squares: impact parameter (b) based centrality



A good agreement with published data.
 $v_2[2]$ and $v_2[4]$ are insensitive to centrality determination method.

The published data taken from: **Niseem Magdy et. al. Published in: Phys.Rev.C 105 (2022) 4, 044901**

Elliptic flow at NICA energies




- v_2 is sensitive to the properties of strongly interacting matter:**



- At $\sqrt{s_{NN}} = 4.5$ GeV pure string/hadronic cascade models (UrQMD, SMASH,...) give similar v_2 signal compared to STAR data
- At $\sqrt{s_{NN}} \geq 7.7$ GeV pure string/hadronic cascade models underestimate v_2 – need hybrid models with QGP phase (vHLE+UrQMD, AMPT with string melting,...)

- Relative v_2 fluctuations ($v_2\{4\}/v_2\{2\}$) observed by STAR experiment can be reproduced both in the string/cascade models (UrQMD, SMASH) and hybrid model (AMPT SM, vHLE+UrQMD)
- Dominant source of v_2 fluctuations: **participant eccentricity fluctuations** in the initial geometry

Elliptic flow at NICA energies




Taranenko et. al.,
Phys. Part. Nuclei **51**, 309–313 (2020)




- **Strong energy dependence of v_2 at $\sqrt{s_{NN}} = 3\text{--}11$ GeV**
 - $v_2 \approx 0$ at $\sqrt{s_{NN}} = 3.3$ GeV and negative below
- **Lack of differential measurements of v_2 at NICA energies (p_T , centrality, PID,...)**
- **v_2 is sensitive to the properties of strongly interacting matter:**
 - At $\sqrt{s_{NN}} = 4.5$ GeV pure string/hadronic cascade models (UrQMD, SMASH,...) give similar v_2 signal compared to STAR data
 - At $\sqrt{s_{NN}} \geq 7.7$ GeV pure string/hadronic cascade models underestimate v_2 – need hybrid models with QGP phase (vHLL+UrQMD, AMPT with string melting,...)

Relative elliptic flow fluctuations






Small value for the $v_2\{4\}/v_2\{2\}$ ratio corresponds to large fluctuation

- Relative v_2 fluctuations ($v_2\{4\}/v_2\{2\}$) observed by STAR experiment can be reproduced both in the string/cascade models (UrQMD, SMASH) and hybrid model (AMPT with string melting, vHLLE+UrQMD)
- Dominant source of v_2 fluctuations: **participant eccentricity fluctuations** in the initial geometry



Elliptic flow at NICA energies



- Relative v_2 fluctuations ($v_2\{4\}/v_2\{2\}$) observed by STAR experiment can be reproduced both in the string/cascade models (UrQMD, SMASH) and hybrid model (AMPT SM, vHLE+UrQMD)
- Dominant source of v_2 fluctuations: **participant eccentricity fluctuations** in the initial geometry

- v_2 is sensitive to the properties of strongly interacting matter:
 - At $\sqrt{s_{NN}} = 4.5$ GeV pure string/hadronic cascade models (UrQMD, SMASH,...) give similar v_2 signal compared to STAR data
 - At $\sqrt{s_{NN}} \geq 7.7$ GeV pure string/hadronic cascade models underestimate v_2 – need hybrid models with QGP phase (vHLE+UrQMD, AMPT with string melting,...)