The 6th international conference on particle physics and astrophysics



Contribution ID : 288

Type : Poster

Linearity of a scintillation crystals response

Thursday, 1 December 2022 13:00 (15)

The results of the study of the specific light output from the energy of X-rays and gamma quanta in the range from 1.5 to 662 keV for scintillation crystals NaI (Tl), CsI (Na), CsI (Tl), Bi4Ge3O12, CdWO4, ZnWO4, Y3Al5O12: Ce, CaF2 (Eu), LSO, YAlO3: Ce. The obtained dependences have a nonlinear form with pronounced minima in the region of the K and L edges of the absorption of the elements that make up the scintillators. When X-ray and gamma quanta are recorded in the energy range from 1 to hundreds of keV, a strong change in the specific ionization losses of the produced photoelectrons dE / dx occurs in the scintillation crystal, there are K and L jumps in the crystal absorption cross section. All this has a significant effect on the light output of scintillators. The possibility of reducing the nonlinearity of the response of scintillators during the detection of gamma radiation is shown.

Primary author(s) : Prof. SALAKHUTDINOV, Gayar (National Research Nuclear University "MEPhI"); Mrs. GRIGORYEVA, Irina (National Research Nuclear University "MEPhI"); Mr. NAUMOV, Peter (National Research Nuclear University "MEPhI")

Presenter(s) : Prof. SALAKHUTDINOV, Gayar (National Research Nuclear University "MEPhI"); Mr. NAU-MOV, Peter (National Research Nuclear University "MEPhI")

Session Classification : Poster Session

Track Classification : Facilities and advanced detector technologies