### The estimation methods of the background induced by the misidentification of a jet as a photon in *pp* collisions at $\sqrt{s}$ = 13 TeV with the ATLAS Detector

K. Kazakova, E. Soldatov, D. Pyatiizbyantseva

on behalf of the ZnunuGamma group



### MEPhl@Atlas meeting 18/11/2022



## **Motivation and goals**

### Motivation:

- To measure the parameters of the Standard Model (SM) to very high precision;
- The search of new physics predicted by the beyond SM (BSM) theories;

Precise measurements of triple and quartic gauge

 couplings sensitive to BSM physics. One of the sensitive processes is Z(vv)γ process.

### <u>Goals:</u>

- To calculate integral and differential in  $E_{
  m T}^{\gamma}$ ,  $N_{
  m jets}$ ,  $p_{
  m T}^{
  m miss}$ ,  $\Delta\phi(\gamma,p_{
  m T}^{
  m miss})$ ,  $p_{
  m T}(Z\gamma)$ ,  $\eta_{\gamma}$ . cross-sections and compare the results with the theory predictions;
- To obtain the strongest up-to-date limits on anomalous neutral triple gauge-boson couplings (aTGCs).





### The backgrounds and the phase space definition

### Signal: Z(vv)y

Backgrounds:

- $\gamma$  + jets via MC  $\rightarrow$  ABCD method based on  $E_{\rm T}^{\rm miss}$  significance and additional variable (or slice method?);
- 26%  $W(\rightarrow lv)\gamma$  fit to data in additional CR based on  $N_{lep}$  (shape from MC);
- 20%  $e \rightarrow \gamma$  fake-rate estimation using Z-peak (tag-n-probe) method;
- 14%  $jet \rightarrow \gamma$  ABCD method based on photon ID and isolation and slice method (shape from Zy QCD);
- 1.9% Z(ll)γ via MC;
- 1.6% ttγ via MC.

FixedCutLoose isolation working point is chosen.

#### Preselections

Preselections	Cut value
$E_{\mathrm{T}}^{\mathrm{miss}}$	$> 130~{ m GeV}$
$E^{\gamma}_{ ext{T}}$	$> 150~{ m GeV}$
Number of photons	$N_\gamma=1$
Lepton veto	$N_e=0,N_\mu=0$

#### **Selections**

Selections	Cut value
$E_{\rm T}^{\rm miss}$ significance	> 11
$ \Delta \phi(E_{ m T}^{ m miss},\gamma) $	> 0.7
$ \Delta \phi(E_{ m T}^{ m miss},j_1) $	> 0.4

# $jet \rightarrow \gamma \ background$

• The background induced by the misidentification of a jet as a photon is studied in this analysis.

**π0 Energy Deposit** 



### ABCD method for *jet* $\rightarrow \gamma$ :

the phase space is splitted into 4 regions based on the

- identification (*tight* or *loose'*) and isolation (*isolated* or *non-isolated*) criteria for photons;
- the main assumption is the absence of correlation between
- identification and isolation criteria.

The estimate of *jet* $\rightarrow \gamma$  events in signal region A derived by ABCD method is 2100  $\pm$  100  $\pm$  300



A large uncertainty is observed. Thus, we have a motivation to estimate  $jet \rightarrow \gamma$  with other methods.

Hadronic jets in which neutral mesons carry a significant fraction of energy may be misidentified as isolated photons.

the SR will be contaminated with  $jet \rightarrow \gamma$ 



More details in back-up

26/10/2022

## Estimation techniques of the slice method I

### <u>Strategy:</u>

- Split the phase space into 4 regions based on kinematic cuts and isolation. The fit region (FR) is a region with relaxed cuts on several variables, where events have a leading photon candidate that is isolated. The SR is a subset of the FR, events in the SR pass all signal kinematic selections.
- 2. The CR1 is a region with relaxed cuts on several variables, where events have a leading photon candidate that is not isolated. The CR2 is a subset of the CR1, events in CR2 pass all signal kinematic selections.
- 3. Photons in all four regions pass the tight selection criteria.



Kinematic cuts

- 4. The fit is performed in the FR, where the *jet*  $\rightarrow \gamma$  process used for the fit is derived from CR1. The relaxed cuts in the FR and the CR1 are applied to dispose of enough statistics.
- 5. Photon is required to pass  $p_T^{\text{cone20}}/p_T^{\gamma} < 0.05$  track isolation in isolated regions. To increase the statistics in non-isolated regions the inverted track isolation  $p_T^{\text{cone20}}/p_T^{\gamma} > 0.05$  is applied.

# Estimation techniques of the slice method II

- 6. The fit can be performed for different variables in the phase-space region with relaxed cuts on these variables.
- 7. To study the dependence of the result on the isolation criteria, control regions CR1 and CR2 are split into successive intervals by the isolation variable, instead of a single integrated anti-isolated region.
- 8. In this way, the number of  $jet \rightarrow \gamma$  background events for a given isolation slice *i* can be estimated as follows:

$$N_{\text{CR1(i)}}^{jet \to \gamma} = N_{\text{CR1(i)}}^{\text{data}} - N_{\text{CR1(i)}}^{Z(\nu\bar{\nu})\gamma} - N_{\text{CR1(i)}}^{\text{bkg}}$$



Kinematic cuts

9. The fit is performed in the FR. Thus, the total number of events in the FR estimated from nonisolated slice of the CR1 is given by:

$$N_{\text{FR}(i)}^{\text{data}} = \alpha \cdot (N_{\text{FR}(i)}^{Z(\nu\bar{\nu})\gamma} + N_{\text{FR}(i)}^{\text{bkg}}) + N_{\text{FR}(i)}^{jet \to \gamma}$$

10. The fitting parameter  $T_{(i)}$  gives the estimated number of  $jet \rightarrow \gamma$  events in the FR:  $N_{FR(i)}^{jet \rightarrow \gamma} \approx T_{(i)} \cdot N_{CR1(i)}^{jet \rightarrow \gamma}$ 

## Estimation techniques of the slice method III

- 11. In this study, a parameter  $\alpha$  is taken to be equal 1. The fit parameter  $T_{(i)}$  is derived for each slice and kinematic variable.
- 12. Finally, the fitted  $jet \rightarrow \gamma$  yield is extrapolated to the signal region. The estimate for each slice and kinematic variable is determined by the equation:

$$N_{\text{SR(i)}}^{jet \to \gamma} = T_{(i)} \cdot (N_{\text{CR2(i)}}^{\text{data}} - N_{\text{CR2(i)}}^{Z(\nu\bar{\nu})\gamma} - N_{\text{CR2(i)}}^{\text{bkg}})$$

FixedCutLoose isolation working point is chosen.
 Isolation working point is defined as:

$$E_{\rm T}^{\rm cone20}/p_{\rm T}^{\gamma} < 0.065$$

Isolation CR1 CR2  $E_{T}^{miss}$  sign. > 0 ET<sup>miss</sup> sign. > 11 ET<sup>miss</sup> > 0 GeV ET<sup>miss</sup> > 130 GeV  $|\Delta \phi (E_T^{miss}, \gamma)| > 0$  $|\Delta \phi (E_T^{miss}, \gamma)| > 0.7$  $|\Delta\phi(E_T^{miss}, j_1)| > 0$  $|\Delta \phi (E_T^{miss}, j_1)| > 0.4$ Tight Tight non-isolated non-isolated SR **CR3 (FR)** ET<sup>miss</sup> sign. > 11  $E_{T}^{miss}$  sign. > 0 ET<sup>miss</sup> > 130 GeV ET<sup>miss</sup> > 0 GeV  $|\Delta \phi (E_T^{miss}, \gamma)| > 0$  $|\Delta \phi (E_T^{miss}, \gamma)| > 0.7$  $|\Delta \phi (E_T^{miss}, j_1)| > 0$  $|\Delta \phi (E_T^{miss}, j_1)| > 0.4$ Tight Tight isolated isolated

Kinematic cuts



Five isolation slices are chosen: [0.065, 0.08, 0.095, 0.115, 0.14]

### Fit process

- The fit was performed for 4 variables:  $E_{\rm T}^{\rm miss}$ ,  $E_{\rm T}^{\rm miss}$  significance,  $|\Delta \phi(\gamma, \vec{p}_{\rm T}^{\rm miss})|$  and  $|\Delta \phi(j_1, \vec{p}_{\rm T}^{\rm miss})|$
- The fitting parameter  $T_{(i)}$  is derived from the fit for each slice and variable.

The results of the fit for slice 1 [0.065, 0.08]:



Pre-fits and post-fits for other slices are performed in back-up

### The results of the fit

#### Result of the fit for $Z\gamma$ QCD Sherpa generator:

	Slice	$T_1$	, $E_{\mathrm{T}}^{\mathrm{miss}}$	$T_2, E_{\mathrm{T}}^{\mathrm{min}}$	<sup>iss</sup> sign	$. \mid T_3, \mid \Delta(E_{\mathrm{T}}^{\mathrm{miss}}, j_1)$	$ T_4,$	$ \Delta(E_{\mathrm{T}}^{\mathrm{miss}},\gamma) $		
	1	3.13	$\frac{1}{1000} \pm 0.07$	3.23 =	E 0.08	$3.06 \pm 0.08$	2	$.71 \pm 0.07$		
	2	3.81	$\pm 0.08$	3.88 =	E 0.09	$3.56 \pm 0.08$	2	$.99\pm0.08$	The f	it
	3	4.00	$0 \pm 0.09$	4.05 =	E 0.09	$3.68\pm0.09$	3	$.01 \pm 0.09$	para	meters $T_{(i)}$
	4	4.96	$5\pm0.11$	5.02 =	E 0.12	$4.47 \pm 0.11$	3	$.42 \pm 0.11$		
	5	1.95	$6 \pm 0.05$	1.96 =	E 0.05	$1.53 \pm 0.04$	1	$.00 \pm 0.04$		
				Slice	Obse	erved $N_{CR2(i)}^{jet \to \gamma}$				
				1	4	$144 \pm 22$				
				2		$320 \pm 19$		Observed in	$et \rightarrow v eve$	nts in
				3		$265 \pm 17$		the CR2		
				4		$207 \pm 15$				
				5		$363 \pm 22$				
$N_{SR}^{jet}$	$E_{(i)}^{t \to \gamma}, E_{\mathrm{T}}^{\mathrm{n}}$	niss	$N_{SR(i)}^{jet \to \gamma},$	$E_{\rm T}^{\rm miss}$ s	ign.	$N_{SR(i)}^{jet \to \gamma},  \Delta(E_{\rm T}^{\rm miss})\rangle$	$, j_1)  $	$N_{SR(i)}^{jet \to \gamma},  \Delta($	$(E_{\mathrm{T}}^{\mathrm{miss}},\gamma) _{-}$	
13	$93 \pm 74$	$4 \mid$	143	$7\pm78$		$1358 \pm 74$		1205 =	± 66	<i>iet</i> $\rightarrow$ <i>v</i> events
12	$17 \pm 70$	6	123	$9\pm78$		$1139 \pm 72$		$956~\pm$	- 62	in the SR for
10	$58 \pm 71$	1	107	$2\pm73$		$972\pm 66$		$795~\pm$	= 56	each slice
10	$26 \pm 78$	8	103	$7\pm79$		$924 \pm 71$		$706 \pm$	= 56	
70	$08 \pm 46$	;	709	$9\pm46$		$556 \pm 37$		$364 \pm$	= 26	

Slice

2

3

4

5

### Linear extrapolation



## The sources of the systematics

### Systematic uncertainties come from:

- The uncertainty in the choice of the extrapolation target for the isolation scan, estimated by changing the isolation target by  $\pm 1\sigma$ ;
- The uncertainty comes from different generators.

Variable	Estimate in $x = 0.002$	Variable	$Z\gamma$ QCD MadGraph
$E_{\mathrm{T}}^{\mathrm{miss}}$	$1920 \pm 110$	$E_{\mathrm{T}}^{\mathrm{miss}}$	$2010 \pm 120$
$E_{\rm T}^{\rm miss}$ sign.	$1990 \pm 120$	$E_{\rm T}^{\rm miss}$ sign.	$2040 \pm 120$
$ \Delta(E_{\rm T}^{\rm miss}, j_1) $	$1970 \pm 110$	$ \Delta(E_{\mathrm{T}}^{\mathrm{miss}},j_{1}) $	$2010 \pm 110$
$ \Delta(E_{\mathrm{T}}^{\mathrm{miss}},\gamma) $	$1820\pm90$	$ \Delta(E_{\mathrm{T}}^{\mathrm{miss}},\gamma) $	$1900 \pm 100$
	ts. δ = 98 events	1982 ev	vents, δ = 169 events

- Total systematic uncertainty is 195 events.
- Thus, the estimate of *jet*  $\rightarrow \gamma$  events in signal region A by slice method is 1810 ± 90 ± 200.
- The estimate of *jet* $\rightarrow \gamma$  events in signal region A derived by ABCD method is 2100  $\pm$  100  $\pm$  300.
- The final estimates for different methods coincide within the uncertainty. W/o  $|\Delta\phi(\gamma, \vec{p}_T^{\text{miss}})|$  the estimate is 1860 ± 70 ± 180 events.

#### MEPhI@Atlas

### Likelihood-based approach I

The main idea: to fit signal and other backgrounds distributions except jet  $\to \gamma$  to data in all ABCD regions

The essence of the method is to perform a fit of the likelihood function, which is defined as:

$$L(N_{ji}|f_{F_{ji}}, f_{N_j}) = \prod_{j=A}^{B,C,D} \prod_{i=1}^{N_{bins}} \text{Pois}(N_{ji}|\nu_{b_{ji}} + \nu_{\gamma_{ji}}f_{F_{ji}} + \nu_{s_{ji}}f_{N_j})$$

where model parameters are defined as:

- $N_{ji}$  the number of the data events in each region and bin;
- $f_{N_i}$  varying parameter for signal in each region;
- $f_{F_{ii}}$  varying parameter for estimated background in each region and bin;
- $\mathcal{V}_{b_{ji}}$  the number of events in MC backgrounds (excl. jet  $\rightarrow \gamma$  );
- $\mathcal{V}_{S_{ji}}$  the number of signal events;
- $\mathcal{V}_{\gamma_{ji}}$  the number of estimated background (jet ightarrow ) events.

#### MEPhl@Atlas

### Likelihood-based approach II

• Likelihood based approach is constructed with the assumption that R = 1 for each bin in the distribution for jet  $\rightarrow \gamma$  background:

$$1 = \frac{\nu_{\gamma_{Ai}} f_{F_{Ai}} \cdot \nu_{\gamma_{Di}} f_{F_{Di}}}{\nu_{\gamma_{Bi}} f_{F_{Bi}} \cdot \nu_{\gamma_{Ci}} f_{F_{Ci}}}$$

To avoid the redundancy of the model the following limitation is applied:

$$f_{F_{Bi}} = f_{F_{Di}}$$

The search of maximum of likelihood function is performed with **RooFit** toolkit:

$$\frac{\partial L}{\partial f_{F_{ji}}} = 0, \quad \frac{\partial L}{\partial f_{N_j}} = 0$$
SR:  $N_{A}^{jet \to \gamma} = \nu_{\gamma} \dots f_{F_{ji}}$ 

This way the number of jet  $\rightarrow \gamma$  events in SR:  $N_A^{\gamma \gamma \gamma} = \nu_{\gamma_{Ai}} f_{F_{Ai}}$ 

The proposed method significantly reduces the number of steps to be done to obtain the estimate compared to ABCD-method

### MC samples

The likelihood-based approach is applied to associated Zy production with Z-boson decaying

• into neutrinos (Z  $\rightarrow$  vv). One of the backgrounds comes from  $\gamma$ +j events. Zj events come from jet  $\rightarrow \gamma$  misidentification

The processes considered in the analysis are generated in

- MadGraph5 MC event generator using pp collisions with  $\sqrt{s}$ 
  - = 13 TeV and the integrated luminosity of 139 fb<sup>-1</sup>
- Pythia8 is used for parton showering and hadronization,
   Delphes is used for detector simulation.

Thus the study uses Asimov data which is not real data but the sum of MC generated processes, the likelihood-based estimate of jet  $\rightarrow \gamma$  background and MC prediction should coincide. It is so-called «closure test».

Selection	Cut value
$E_{\mathrm{T}}^{\mathrm{miss}}$	> 130 GeV
$E_{\mathrm{T}}^{\gamma}$	> 150  GeV
Number of tight photons	$N_{\gamma} = 1$
Lepton veto	$N_e=0,N_\mu=0$

Event selection criteria for Zγ candidate events

# The results of the fit

100

### The fit was performed for $\phi_{\gamma}$ and $\eta_{\gamma}$ :

The final estimate is chosen based on the  $\chi^2/N_{d.o.f.}$ value in the SR and R-factor.

NL.		$\phi_\gamma$		$\eta_{\gamma}$			
<sup>1</sup> <b>v</b> bins	Estimate	R-factor	$\chi^2/N_{d.o.f.}$	Estimate	R-factor	$\chi^2/N_{d.o.f.}$	
6	$3255^{+111}_{-106}$	$1.04\pm0.03$	0.45	$3238^{+129}_{-125}$	$1.03\pm0.03$	0.39	
7	$2906^{+110}_{-108}$	$0.94\pm0.03$	0.73	$3243^{+126}_{-122}$	$1.04\pm0.02$	0.55	
8	$3179^{+117}_{-108}$	$1.04\pm0.03$	0.73	$3276^{+141}_{-137}$	$1.04\pm0.02$	0.26	
9	$3119^{+130}_{-127}$	$1.01\pm0.03$	0.62	$3251^{+133}_{-130}$	$1.05\pm0.02$	0.50	

The systematic uncertainties were derived by

- variating the value of isolation gap by  $\pm \sigma$  in nonisolated control regions.
- The estimate of jet  $\rightarrow \gamma$  events in SR obtained by ¢ ٍ, [Rad] likelihood method is  $N_A^{jet \rightarrow \gamma} = 3179^{+117}_{-108} \pm 69$  for  $\phi_{\gamma}$  and  $N_A^{jet \rightarrow \gamma} = 3243^{+126}_{-122} \pm 48$  for  $\eta_{\gamma}$
- The MC prediction is  $N_A^{jet \rightarrow \gamma} = 3093 \pm 178$  events



# Summary

The estimate of *jet* $\rightarrow \gamma$  events in signal region A is derived by ABCD method. The estimate is 2100  $\pm$  100  $\pm$  300 events.

The alternative slice method is performed for  $jet \rightarrow \gamma$  estimation process. The estimate of  $jet \rightarrow \gamma$  events in signal region A derived by slice method is 1810 ± 90 ± 200. The final estimates for the methods coincide within the uncertainty.

The alternative likelihood-based method of estimation of  $jet \rightarrow \gamma$  events was developed. It uses the information about the shape of the distributions in the regions and provides a much simpler way to obtain the estimate of the number of background events.

### Thank you for your attention!



# R factor Zj and W(τν) in MC

Working point	loose'2	loose'3	loose'4	loose'5
FCTight	$1.06 \pm 0.16$	$1.15 \pm 0.16$	$1.17 \pm 0.15$	$1.30 \pm 0.17$
FCTight (inversion)	$1.05 \pm 0.14$	$1.21 \pm 0.15$	$1.38 \pm 0.16$	$1.65 \pm 0.19$
FCTCaloOnly	$1.18 \pm 0.13$	$1.31 \pm 0.13$	$1.37 \pm 0.13$	$1.54 \pm 0.14$
FCLoose	$1.0 \pm 0.2$	$1.0 \pm 0.2$	$1.0 \pm 0.2$	$1.3 \pm 0.2$
FCLoose (inversion)	$1.11 \pm 0.13$	$1.23 \pm 0.12$	$1.34 \pm 0.12$	$1.60 \pm 0.13$



$$E_{\rm T}^{\rm miss} \text{ significance } =$$
$$|\vec{E}_{\rm T}^{\rm miss}|^2 / (\sigma_{\rm L}^2 (1 - \rho_{\rm LT}^2))$$

 $\sigma_{\rm L}$  is the total variance in the longitudinal direction to the  $E_{\rm T}^{\rm miss}$  $ho_{\rm LT}$  is the correlation factor of the longitudinal L and transverse T measurement

### Isolation distributions (loose'3)

- Tight(MC)

Tight(MC)

- Loose'3(MC)

Loose'3(Data)

----

20

Loose'3(MC)

- Loose'3(Data)





K. Kazakova (NRNU MEPhl)

MEPhI@Atlas

## R factor in data

MEPhI@Atlas

	<b>FixedCutLoose (inverted)</b> , w/o upper cut							
			MC					
		loose'2	loose'3	loose'4	loose'5			
F	R-factor	$1.11 \pm 0.13$	$1.23 \pm 0.12$	$1.34 \pm 0.12$	$1.60 \pm 0.13$			
Data-driven								
	Cut	loose'2	loose'3	loose'4	loose'5			
	4.5	$0.97\pm0.10$	$1.05 \pm 0.10$	$1.05 \pm 0.09$	$1.06 \pm 0.08$			
	4.6	$1.00 \pm 0.10$	$1.08 \pm 0.10$	$1.06 \pm 0.09$	$1.07 \pm 0.08$			
	4.75	$1.03 \pm 0.10$	$1.05 \pm 0.10$	$1.07 \pm 0.09$	$1.09 \pm 0.08$			
	9.5	$1.04 \pm 0.09$	$1.03 \pm 0.08$	$0.98 \pm 0.07$	$0.97 \pm 0.07$			
	10.0	$1.04 \pm 0.09$	$1.03 \pm 0.08$	$0.98 \pm 0.07$	$0.98 \pm 0.07$			
	10.5	$1.02 \pm 0.09$	$1.02 \pm 0.08$	$0.95 \pm 0.07$	$0.96 \pm 0.07$			
	11.0	$1.06 \pm 0.09$	$1.02 \pm 0.08$	$0.97 \pm 0.07$	$0.96 \pm 0.07$			
		<b>EiredCut</b>	mbt Cala Oraliz					
	FixedCutTightCaloOnly, w/o upper cut							
		loogo?2		looso'4	loogo?5			
	D	100se 2	100se 5	100se 4	IOOSE 5			
	R-factor	$1.18 \pm 0.13$	$1.31 \pm 0.13$	$1.37 \pm 0.13$	$1.54 \pm 0.14$			
			Data-driver	1				
	Cut	loose'2	loose'3	loose'4	loose'5			
	9.45	$1.15 \pm 0.07$	$1.21 \pm 0.06$	$1.20 \pm 0.06$	$1.23 \pm 0.06$			
(	9.95	$1.14 \pm 0.06$	$1.20 \pm 0.06$	$1.19 \pm 0.06$	$1.22 \pm 0.06$			
	10.45	$1.15 \pm 0.06$	$1.20 \pm 0.06$	$1.19 \pm 0.05$	$1.21 \pm 0.05$			
	$\overline{10.45}$	$1.21 \pm 0.07$	$1.26 \pm 0.06$	$1.24 \pm 0.06$	$1.26 \pm 0.06$			

F	<b>FixedCutTight</b> (inverted), w/o upper cut							
		MC						
	loose'2	loose'3	loose'4	loose'5				
R-factor	$1.05 \pm 0.14$	$1.21 \pm 0.15$	$1.38 \pm 0.16$	$1.65 \pm 0.19$				
	Data-driven							
Cut	loose'2	loose'3	loose'4	loose'5				
9.45	$1.10 \pm 0.08$	$1.15 \pm 0.07$	$1.11 \pm 0.06$	$1.16 \pm 0.06$				
9.95	$1.09 \pm 0.07$	$1.15 \pm 0.07$	$1.12 \pm 0.06$	$1.16 \pm 0.06$				
10.20	$1.08 \pm 0.07$	$1.14 \pm 0.07$	$1.11 \pm 0.06$	$1.15 \pm 0.06$				
10.45	$1.10 \pm 0.07$	$1.15 \pm 0.07$	$1.13 \pm 0.06$	$1.17 \pm 0.06$				
FivedCutTight_upper_gut = 25.45 CeV								
	Incucuti	MC	ut – 20.40 C					
	loose'2	loose'3	loose'4	loose'5				
R-factor	$1.07 \pm 0.16$	$1.17 \pm 0.17$	$1.18 \pm 0.16$	$1.31 \pm 0.17$				
	I	Data-driver	1					
Cut	loose'2	loose'3	loose'4	loose'5				
8.45	$1.15 \pm 0.13$	$1.16 \pm 0.12$	$1.16 \pm 0.11$	$1.21 \pm 0.11$				
8.95	$1.11 \pm 0.13$	$1.11 \pm 0.12$	$1.14 \pm 0.11$	$1.17 \pm 0.11$				
9.45	$1.19 \pm 0.14$	$1.22 \pm 0.13$	$1.27 \pm 0.13$	$1.30 \pm 0.12$				
9.95	$1.16 \pm 0.14$	$1.17 \pm 0.13$	$1.23 \pm 0.12$	$1.28 \pm 0.12$				
10.45	$1.19 \pm 0.14$	$1.20 \pm 0.14$	$1.22 \pm 0.12$	$1.26 \pm 0.12$				
FixedCu	tLoose was cł	nosen. In orde	r to decrease	syst.				

uncert. the loose'3 was chosen

#### K. Kazakova (NRNU MEPhl)

# jet $\rightarrow \gamma$ background estimation (loose'3)

Event yields for the data and non-jet  $\rightarrow \gamma$  background processes considered in the ABCD method

	Data	$W\gamma \ QCD$	$W\gamma EWK$	$e\to\gamma$	$tt\gamma$	$\gamma+{ m jet}$	$Z(ll)\gamma$
А	$26523 \pm 163$	$3936 \pm 23$	$136.3 \pm 0.7$	$3039 \pm 12$	$234 \pm 3$	$5262 \pm 53$	$285\pm5$
В	$1475 \pm 38$	$52 \pm 4$	$1.86 \pm 0.08$	$8.95\pm0.03$	$1.3 \pm 0.2$	$0.6 \pm 0.4$	$1.0 \pm 0.6$
С	$2568 \pm 51$	$60 \pm 2$	$2.16 \pm 0.09$	$61.4 \pm 0.2$	$4.2\pm0.4$	$76 \pm 6$	$4.8\pm0.5$
D	$1443 \pm 38$	$2.7 \pm 0.6$	$0.17\pm0.02$	$0.0715 \pm 0.0002$	$0.35 \pm 0.13$	$0\pm 0$	$0\pm 0$

$$N_{\rm A}^{\rm sig} = \widetilde{N}_{\rm A} - R(\widetilde{N}_{\rm B} - c_{\rm B}N_{\rm A}^{\rm sig}) \frac{\widetilde{N}_{\rm C} - c_{\rm C}N_{\rm A}^{\rm sig}}{\widetilde{N}_{\rm D} - c_{\rm D}N_{\rm A}^{\rm sig}}$$
$$N_{\rm A}^{\rm sig} = \frac{b - \sqrt{b^2 - 4ac}}{2a}$$

he	signal	leakage	parameters
IIC	Signat	icanage	parameters

_

5:	Event yields signal:					
	$Z(\nu\bar{\nu})\gamma$ QCD	$Z(\nu\bar{\nu})\gamma \ \mathrm{EWK}$				
А	$10513 \pm 8$	$152.1 \pm 0.3$				
В	$98.0\pm0.8$	$2.14 \pm 0.04$				
С	$161.5 \pm 1.0$	$2.31 \pm 0.04$				
D	$5.3 \pm 0.2$	$0.135 \pm 0.009$				

$$\begin{aligned} a &= c_D - Rc_B c_C; \\ b &= \widetilde{N}_{\rm D} + c_D \widetilde{N}_{\rm A} - R(c_B \widetilde{N}_{\rm C} + c_C \widetilde{N}_{\rm B}); & \text{With F} \\ c &= \widetilde{N}_{\rm D} \widetilde{N}_{\rm A} - R \widetilde{N}_{\rm C} \widetilde{N}_{\rm B}. \end{aligned}$$

h R by data-driven 📃

$$N_A^{jet
ightarrow\gamma}=$$
 2078<sup>+100</sup><sub>-97</sub>

# Systematic uncertainty I

### Systematic uncertainties come from:

- non-tight definition and isolation gap choice. Variation for ±1σ changes in data yield
- different generators
- imperfect photon iso/ID modeling

Dittoront		nrimo	and	ICOL	ation	C D	n
	<b>LUUJE</b>		anu	1301	αιισι	ya	μ

Central value (with $R_{data}$ )	2078
loose'2	+327
loose'4	-111
loose'5	-173
Iso gap $+0.25$ GeV	+48
Iso gap $-0.35$ GeV	+29

 $R_{\text{data}}^{\text{iso gap } +0.25 \text{ GeV}} = 1.07 \pm 0.11$  $R_{\text{data}}^{\text{iso gap } -0.35 \text{ GeV}} = 1.06 \pm 0.09$ 

Iso gap, GeV	$N_B$	$N_D$
-0.40	$1524 \pm 39$	$1488 \pm 39$
-0.35	$1518 \pm 39$	$1482 \pm 38$
-0.30	$1513 \pm 39$	$1477 \pm 38$
-0.25	$1503 \pm 39$	$1474 \pm 38$
-0.20	$1497 \pm 39$	$1468 \pm 38$
2.0	$1475 \pm 38$	$1443 \pm 38$
+0.15	$1448 \pm 38$	$1416 \pm 38$
+0.20	$1443 \pm 38$	$1404 \pm 37$
+0.25	$1437 \pm 38$	$1398 \pm 37$

δ = 16%

The choice of loose prime 3 reduced the systematic uncertainty from 32% to 16%

MEPhI@Atlas

# Systematic uncertainty II

#### **Different generators:**

	Different generators					
Signal leakage parameters	MadGraph+Pythia8, Sherpa 2.2	MadGraph+Pythia8, MadGraph+Pythia8	Relative deviation			
c <sub>B</sub>	$0.00939 \pm 0.00007$	$0.0155 \pm 0.0004$	39%			
$\mathrm{c}_C$	$0.01536 \pm 0.00010$	$0.0156 \pm 0.0004$	1.5%			
$c_D$	$0.00051 \pm 0.000028$	$0.00077 \pm 0.00009$	34%			
$jet \rightarrow \gamma$ est. (with $R_{data}$ )	2078	2061	0.8%			

• Uncertainty coming from signal leakage is obtained  $\delta = 0.8\%$ 

 $R_{\rm data}^{\rm diff.gen.} = 1.10 \pm 0.10$ 

Systematic uncertainty come from imperfect photon iso/ID modeling:

• 
$$\sigma_{iso}^{c_B} = \delta_{iso}^{eff} \cdot (c_B + 1)/c_B$$
  
•  $\sigma_{ID}^{c_C} = \delta_{ID}^{eff} \cdot (c_C + 1)/c_C$   
•  $\sigma_{iso}^{c_D} = \delta_{iso}^{eff} \cdot (c_B + 1)/c_B$   
•  $\sigma_{ID}^{c_D} = \delta_{ID}^{eff} \cdot (c_C + 1)/c_C$   
•  $\sigma_{ID}^{c_D} = \delta_{ID}^{eff} \cdot (c_C + 1)/c_C$ 

Total systematics: δ<sub>Data</sub> = 16%







E<sup>miss</sup> significance





### The results of the fit

• Events in FR:

Data	Background (excl. $jet \to \gamma$ )	Signal
$93513 \pm 306$	$65902 \pm 1019$	$13742 \pm 13$

Slice	Data	Background (excl. $jet \to \gamma$ )	Signal (Sherpa)	Signal (MadGraph)
1	$4572 \pm 68$	$78 \pm 5$	$20.7 \pm 0.4$	$27 \pm 2$
2	$3776 \pm 61$	$48 \pm 4$	$18.4\pm0.3$	$20 \pm 2$
3	$3642 \pm 60$	$79 \pm 4$	$20.1\pm0.3$	$26 \pm 2$
4	$2916 \pm 54$	$41 \pm 3$	$21.0\pm0.3$	$27 \pm 2$
5	$7672 \pm 88$	$241 \pm 6$	$128.1\pm0.8$	$153 \pm 5$

#### Events in CR1 for each slice:

Slice	Data	Background (excl. $jet \to \gamma$ )	Signal (Sherpa)	Signal (MadGraph)
1	$463 \pm 22$	$8.4 \pm 0.9$	$10.1 \pm 0.3$	$16.7 \pm 1.3$
2	$337 \pm 18$	$8 \pm 3$	$9.3\pm0.3$	$12.7 \pm 1.2$
3	$286 \pm 17$	$11.6 \pm 0.9$	$9.7\pm0.2$	$15.5 \pm 1.3$
4	$223 \pm 15$	$5.5 \pm 0.9$	$10.9\pm0.3$	$18.6 \pm 1.3$
5	$471 \pm 22$	$41 \pm 2$	$67.0\pm0.6$	$105 \pm 3$

Events in CR2 for each slice:

### The results of the fit

#### Result of the fit for Zy QCD MadGraph:

	Slice	$T_1, E_{\mathrm{T}}^{\mathrm{miss}}$	$T_2, E_{\mathrm{T}}^{\mathrm{miss}}$ sign	n. /	$T_3,  \Delta(E_{\rm T}^{\rm min})\rangle$	$ iss, j_1) $	$T_4,  _4$	$\Delta(E_{\mathrm{T}}^{\mathrm{miss}},\gamma) $	_
	1	$3.38 \pm 0.07$	$3.42 \pm 0.08$	3	$3.26 \pm 0$	0.08	2.9	$96 \pm 0.07$	_
	2	$4.09 \pm 0.08$	$4.09 \pm 0.09$	)	$3.78 \pm 0$	0.08	3.2	$28\pm0.08$	
	3	$4.32 \pm 0.10$	$4.27 \pm 0.10$	)	$3.91 \pm 0$	0.09	3.3	$30 \pm 0.09$	
	4	$5.35 \pm 0.12$	$5.29 \pm 0.12$	2	$4.76 \pm 0$	0.11	3.7	$76 \pm 0.11$	
	5	$2.12 \pm 0.05$	$2.06 \pm 0.05$	5	$1.63 \pm 0$	0.04	1.1	$11 \pm 0.04$	
et 2200 2000 1800 1600 1400 1200 1000 800 600 400	ATLAS √s=13 Te	S Internal	• Fit to $E_T^{miss}$ signif. • Fit to $ \Delta\phi(E_T^{miss}, j) $ • Fit to $ \Delta\phi(E_T^{miss}, \gamma) $ • Fit to $E_T^{miss}$		Slice 1 2 3 4 5	Obse 4 3 2 2 3	erved 44 ± 820 ± 265 ± 207 ± 863 ±	$     \begin{array}{r} N_{CR2(i)}^{jet \to \gamma} \\ = 22 \\ = 19 \\ = 17 \\ = 15 \\ = 22 \end{array} $	Obsei event
C	0.02 0.04	0.06 0.08 0.	1 0.12 0.14						
			E <sup>cone20</sup> /p <sub>T</sub>						
Slice	$N_{SR(i)}^{jet \to \gamma}, E_{\mathrm{T}}^{\mathrm{m}}$	iss $N_{SR(i)}^{jet \to \gamma}$ ,	$E_{\rm T}^{\rm miss}$ sign.	$N_S^{j}$	$S_{R(i)}^{iet \to \gamma},  \Delta($	$(E_{\mathrm{T}}^{\mathrm{miss}}, E_{\mathrm{T}})$	$j_1) $	$N_{SR(i)}^{jet \to \gamma},  \Delta$	$\Delta(E_{\mathrm{T}}^{\mathrm{miss}},\gamma) $
1	$1482 \pm 79$	) 149	$7 \pm 81$		1427 =	$\pm 78$		1296	$\pm$ 71
2	$1294 \pm 81$	129	$3\pm82$		1196 =	$\pm 75$		1036	$\pm 67$
3	$  1118 \pm 78$	3   110	$5\pm77$		1010 =	$\pm 70$		854	$\pm 61$
4	$1065 \pm 84$	l   105	$3\pm 83$		$947~\pm$	= 75		749	$\pm 61$
5	$689 \pm 49$	670	$0 \pm 48$		$529 \pm$	= 38		362	$\pm 28$

The fit parameter  $T_{(i)}$ 

Observed  $jet \rightarrow \gamma$  events in the CR2

 $jet \rightarrow \gamma$  events in the SR for each slice

### **Distributions**



 $\Delta \phi(\gamma, MET)$  is not sensitive to the jet  $\rightarrow \gamma$  background then estimating with slice method?

K. Kazakova (NRNU MEPhl)

MEPhI@Atlas

### The results of the fit



### The results of the fit

