



Contribution ID : 607

Type : Oral talk

## On generalized Melvin solutions for Lie algebras of rank 4

*Tuesday, 6 October 2020 11:00 (20)*

We consider generalized Melvin-like solutions associated with Lie algebras of rank 4 (namely,  $A_4, B_4, C_4, D_4$ , and the exceptional algebra  $F_4$ ) corresponding to certain internal symmetries of the solutions. The system under consideration is a static cylindrically-symmetric gravitational configuration in  $D$  dimensions in presence of four Abelian 2-forms and four scalar fields. The solution is governed by four moduli functions  $H_s(z)$  ( $s = 1, \dots, 4$ ) of squared radial coordinate  $z = \rho^2$  obeying four differential equations of the Toda chain type. These functions turn out to be polynomials of powers  $(n_1, n_2, n_3, n_4) = (4, 6, 6, 4), (8, 14, 18, 10), (7, 12, 15, 16), (6, 10, 6, 6), (22, 42, 30, 16)$  for Lie algebras  $A_4, B_4, C_4, D_4, F_4$ , respectively. The asymptotic behaviour for the polynomials at large distances is governed by some integer-valued  $4 \times 4$  matrix  $\nu$  connected in a certain way with the inverse Cartan matrix of the Lie algebra and (in  $A_4$  case) the matrix representing a generator of the  $Z_2$ -group of symmetry of the Dynkin diagram. The symmetry properties and duality identities for polynomials are obtained, as well as asymptotic relations for solutions at large distances. We also calculate 2-form flux integrals over 2-dimensional discs and corresponding Wilson loop factors over their boundaries.

**Primary author(s)** : Dr. BOLOKHOV, Sergei

**Co-author(s)** : Dr. IVASHCHUK, Vladimir

**Presenter(s)** : Dr. BOLOKHOV, Sergei

**Session Classification** : Gravitation and Cosmology

**Track Classification** : Gravitation and cosmology