



Contribution ID : 693

Type : Oral talk

The Nambu Sum Rule in the Composite Two Higgs Doublet Model

Tuesday, 6 October 2020 17:05 (15)

The spectrum of spinless modes in a model with the $SU(2)_L \times SU(2)_R$ symmetrical four-quark interaction proposed by Miransky, Tanabashi, and Yamawaki is studied. For the sake of simplicity, only four-fermion interactions of top and bottom quarks are considered. The spinless modes result from spontaneous electroweak interaction symmetry breaking and are coupled quark–antiquark states associated with $SU(2)$ two Higgs doublets. Their dynamics is described by the effective Lagrangian obtained by the Schwinger–DeWitt method. The spectrum is represented by excitations of five types, the mass of each being expressed by the parameters of the model. It is shown that the model yields phenomenologically acceptable values of both the mass of quarks $m_t=173$ GeV and $m_b=4.18$ GeV and the mass of the standard Higgs state $m(\chi_{-1})=125$ GeV. The masses of the particles that comprise the second Higgs doublet $m(\hat{h}^\mp)=275$ GeV, $m(\chi_{-1})=325$ GeV and $m(\phi_{-0})=125$ GeV, have been calculated. The Nambu sum rule and the conditions for satisfying it in the theories with the broken $U(1)_A$ symmetry are discussed.

Primary author(s) : Prof. OSIPOV, Alexander (Joint Institute for Nuclear Research, Dubna, Moscow oblast, 141980 Russia); Mr. KHALIFA, Mahmoud (Moscow Institute of Physics and Technology, Dolgoprudnyi, Moscow oblast, 141701 Russia, Department of Physics, Al-Azhar University, Cairo, 11751 Egypt)

Presenter(s) : Mr. KHALIFA, Mahmoud (Moscow Institute of Physics and Technology, Dolgoprudnyi, Moscow oblast, 141701 Russia, Department of Physics, Al-Azhar University, Cairo, 11751 Egypt)

Session Classification : HEP theory

Track Classification : HEP theory