

# **Evolution of anisotropic flow of produced particles from Au+Au collisions at $\sqrt{s} = 7.7 - 200$ GeV in a hybrid models**

A. Demanov, A. Povarov, P. Parfenov, G. Nigmatkulov, A. Taranenko  
(NRNU MEPhI)


5th International Conference on Particle Physics and  
Astrophysics (ICCPA 2020), October 5-9, 2020, Moscow

*This work is supported by: the RFBR according to the research project No. 18-02-40086 and by  
Ministry of Science and Higher Education of the Russian Federation, No 0723-2020-0041.*


# OUTLINE

1. Why measure anisotropic flow?
2. Anisotropic flow ( $V_n$ ) and sQGP at RHIC/LHC
3. Elliptic flow results from Beam Energy Scan (RHIC) and comparison with hybrid models
4. Outlook for flow measurements at NICA

# Phase Diagram of the Strongly-Interacting Matter




**Top RHIC/LHC: validation of the cross over transition leading to the sQGP**




- **Top RHIC energy /LHC - access to high  $T$  and small  $\mu_B$**
- **RHIC-BES/SPS/NICA/FAIR - access to different systems and a broad domain of the  $(\mu_B, T)$ -plane**

$$\frac{\eta}{s}(T, \mu), \frac{\zeta}{s}(T, \mu), c_s(T), \hat{q}(T), \alpha_s(T), \text{ etc}$$



# Anisotropic Collective Flow at RHIC/LHC



Initial eccentricity (and its attendant fluctuations),  $\epsilon_n$ , drives momentum anisotropy,  $v_n$ , with specific viscous modulation



# Anisotropic Collective Flow at top RHIC / LHC



$V_n$  ( $p_T$ , centrality) - sensitive to the early stages of collision.

Important constraint for transport properties: EOS,  $\eta/s$ ,  $\zeta/s$ , etc.

$v_n$  of identified hadrons:

**Mass ordering at  $p_T < 2 \text{ GeV/c}$**

(hydrodynamic flow, hadron rescattering)

**Baryon/meson grouping at  $p_T > 2 \text{ GeV/c}$**

(recombination/coalescence),

Number of constituent quark (NCQ) scaling

**No difference** between particles and antiparticles



Gale, Jeon, et al., Phys. Rev. Lett. 110, 012302

STAR PRL 118 (2017) 212301

# Beam-Energy Dependence of Elliptic Flow ( $v_2$ )

STAR: *Phys. Rev. C* 93 (2016) 14907

*Phys. Rev. Lett.* 110, 142301 (2013)



- Small change in  $v_2(p_T)$  for inclusive and identified charged hadrons (pions, kaons and (anti)protons) as the  $\sqrt{s_{NN}}$  changes by a factor  $\sim 25$  (from 7.7 GeV to 200 GeV).
- Substantial particle-antiparticle split at lower energies

## Goal of this work:

- 1) Perform simulations with hybrid models (vHLL+UrQMD and AMPT), analyse them as in the real experiment and make comparison with RHIC BES published measurements of  $v_2$ .
- 2) Make projections for measurements at NICA energies ( $\sqrt{s_{NN}} = 4 - 11$  GeV)?

# Hybrid models for anisotropic flow at RHIC/LHC

## 1) UrQMD + 3D viscous hydro model vHLLE + UrQMD

Iurii Karpenko, *Comput. Phys. Commun.* 185 (2014), 3016


<https://github.com/yukarpenko/vhllle>

Parameters: from Iu. A. Karpenko, P. Huovinen, H. Petersen, M. Bleicher, *Phys. Rev. C* 91 (2015) no.6, 064901 – good description of STAR BES results for  $v_2$  of inclusive charged hadrons (7.7 – 62.4 GeV)

Initial conditions: model UrQMD

QGP phase: 3D viscous hydro (vHLLE) with crossover EOS (XPT)

Hadronic phase: model UrQMD



## 2) A Multi-Phase Transport model (AMPT) for high-energy nuclear collisions. *The main source codes: Zi-Wei Lin*


(<http://myweb.ecu.edu/linz/ampt/v1.26t9b/v2.26t9b>)

Initial conditions: model HIJING


QGP phase: Zhang's parton cascade for modeling partonic scatterings

Hadronic phase: model ART

Z.W. Lin, C. M. Ko, B.A. Li, B. Zhang and S. Pal:  
*Physical Review C* 72, 064901 (2005).



# Analysis method: Event plane method ( $\eta$ -sub)





TPC (E) half ( $\eta < 0$ )  $\rightarrow \eta_-$   
 TPC (W) half ( $\eta > 0$ )  $\rightarrow \eta_+$

$$v_n = \frac{\langle \cos[n(\varphi_{\eta_{\pm}} - \Psi_{n,\eta_{\mp}})] \rangle}{\sqrt{\langle \cos[n(\Psi_{n,\eta_+} - \Psi_{n,\eta_-})] \rangle}}$$

resolution





STAR Phys. Rev. C 88 (2013) 14902



**The resulting values for event plane resolution for simulated events from hybrid models: vHLE+UrQMD and AMPT are close to STAR experimental data.**





# vHLLE+UrQMD: Elliptic flow at top RHIC energy : $\sqrt{s_{NN}} = 200$ GeV





Reasonable agreement between results of vHLLE+UrQMD model and published PHENIX data for 200 GeV including K<sub>ET</sub>/n<sub>q</sub> scaling

# $V_2$ of charged mesons at RHIC BES ( $\sqrt{s_{NN}} = 27$ GeV)




STAR data: *Phys. Rev. C* 93 (2016) 14907



# $v_2$ of (anti)protons at RHIC BES ( $\sqrt{s_{NN}} = 27$ GeV)


STAR data: *Phys. Rev. C* 93 (2016) 14907



Difference between results from vHLE+UrQMD model and data for protons and antiprotons  
Model predicts that  $v_2$  (protons) <  $v_2$  (antiprotons), data show  $v_2$  (protons) >  $v_2$  (antiprotons)  
and the difference is growing with decreasing of collision energy.

# AMPT: $v_2$ of identified hadrons at RHIC BES ( $\sqrt{s_{NN}} = 27$ GeV)

STAR data: *Phys. Rev. C* 93 (2016) 14907






Difference between results from AMPT model SM and data for all particles – tuning of parameters?

Model also predicts that  $v_2$  (protons) <  $v_2$  (antiprotons) ,  
data show  $v_2$  (protons) >  $v_2$  (antiprotons)


# $v_2$ of protons and antiprotons at RHIC BES ( $\sqrt{s_{NN}} = 11.5$ GeV)

STAR data: *Phys. Rev. C* 93 (2016) 14907



Models AMPT and vHLE+UrQMD predicts also predicts that  $v_2$  (protons) <  $v_2$  (antiprotons), data show  $v_2$  (protons) >  $v_2$  (antiprotons)

# Elliptic flow: Models vs Data comparison for NICA energy range



Pure String/Hadronic Cascade models (no QGP phase) give smaller  $v_2$  signal compared to STAR data for Au+Au  $\sqrt{s_{NN}}=7.7-11.5$  GeV and models give similar  $v_2$  signal compared to STAR data for Au+Au  $\sqrt{s_{NN}}=4.5$  GeV.

# Summary

- We performed a high statistics simulations with hybrid models (vHLL+UrQMD and AMPT) for several points in collision energy from RHIC BES program.
- The events were analysed in a similar way as the real experimental data and results were compared with STAR published results of  $v_2$  for charged pions, kaons and (anti)protons.
- The results from vHLL+UrQMD model are in a better agreement with experimental data than for AMPT (tunning of the input parameters?)  
Both models in the present configuration fails to reproduce the difference between elliptic flow signal of particles and antiparticles: models predict that  $v_2$  (protons) <  $v_2$  (antiprotons) , data show  $v_2$  (protons) >  $v_2$  (antiprotons)

Model/Data comparison for NICA energy range ( 4-11 GeV):

Pure String/Hadronic Cascade models (no QGP phase) give smaller  $v_2$  signal compared to STAR data for Au+Au  $\sqrt{s_{NN}}=7.7-11.5$  GeV and models give similar  $v_2$  signal compared to STAR data for Au+Au  $\sqrt{s_{NN}}=4.5$  GeV.

Thank you for your attention!

# Anisotropic Flow in Heavy-Ion Collisions: 1988

**Provides reliable estimates of pressure & pressure gradients**


**Can address questions related to thermalization**

**Gives insights on the transverse dynamics of the Medium**

**Provides access to the transport properties of the medium: EOS, sound speed ( $c_s$ ), viscosity, etc**

*Plastic Ball Collaboration,  
H.H. Gutbrod et al., Phys. Lett. B216, 267 (1989)*

*Fourier Expansion for azimuthal anisotropy,  
Cheuk-Yin WONG, Physics Letters, 88B, p 39 (1979)*






# Excitation function of differential elliptic flow

EPJ Web Conf. 204 (2019) 03009

FOPI (15-29%)  
E895 (12-25%)  
STAR (10-40%)



High precision differential measurements of anisotropic flow? 17