



Approaches in centrality measurements of heavy ion collisions with forward calorimeters at MPD/NICA facility



Volkov Vadim

09/10/2020

Overview

- MPD@NICA facility
- FHCAL@MPD and energy depositions for LAQGSM, DCM-SMM models.
- A few methods for centrality determination:
 - a) Correlations of transverse and longitudinal energy components,
 - b) 2D-fit of FHCAL energy distributions.
- Further improvement of the centrality determination
 - a) Subtraction of pion energy contamination and evaluation of spectator's energy.
- Simulations are made for LAQGSM and DCM-SMM fragmentation models for Au-Au collisions with $\sqrt{S_{NN}} = 11 \text{ GeV}$ energy.

NICA facility




Energy range: $\sqrt{s_{NN}}=4-11$ GeV (Au-beam)


Detector requirements:

- Large homogenous acceptance : 2π in azimuthal angle, $|\eta| < 3$, $0.1 < p_t < 3$ GeV/c
- High efficient 3-D track reconstruction
- Powerful PID: p/K up to 1.5 GeV/c, K/p up to 3.0 GeV/c
- ECAL for g , e
- Careful event characterization: impact parameter & event plane reconstruction
- High event rate capability up to ~ 6 kHz

- tracking:
up to $|\eta| < 1.8$ (TPC)
- PID:
had., e , γ (TOF, TPC, ECAL)
- Reaction:
centrality & plane determination (FHCAL)




FHCal@MPD



Two upstream/downstream parts

44 individual modules




Beam hole


FHCAL modules

Non-spectator's contributions

- The main purpose of the FHCal is to detect spectators and to provide an experimental measurement of a heavy-ion collision centrality and orientation of its reaction plane.
- There is an ambiguity in FHCal energy deposition for central/peripheral events due to the fragments (bound spectators) leak into beam hole.
- FHCal measures not only spectator's but also pion's energies.




Energy depositions in FHCaI for different models



- Energy depositions are quite different for different fragmentation models.
- Results would depend on the fragmentation model.
- FHCaI detects not only the spectators but also the produced particles and wounded nucleons from participant region.

Transverse energy distributions are wider for central events and narrower for the peripheral collisions.



This feature can be used for the separation of central/peripheral events.

Correlation between transverse and longitudinal energies in FHCaI

- LAQGSM and DCM-SMM models for $\sqrt{S} = 11$ AGeV are used.

- The E_T and E_L energies are transverse and longitudinal energies respectively.





$$E_T = \sum E_i \sin\theta_i, E_L = \sum E_i \cos\theta_i$$

- The $(E_T - E_L)$ histograms are divided into ten parts, 10% of events in each part, 10%-clusters are separated from one another by perpendiculars to the envelope.

- b-distributions for each centrality bin are fitted by Gauss.


- The separation of central and peripheral events with DCM-SMM model is clearly worse.

New approaches are needed



2D-linear fit method (linear approach)




Energy distribution in FHCAL modules



- In this method the space energy distribution in FHCAL modules is used.
- The energy in the histogram is uniformly distributed in FHCAL modules according to the polar angle.
- The histogram is fitted by a symmetrical cone (linear approximation).
- Weight of each bin is proportional of the energy deposited in corresponding FHCAL module.
- This fit provides the new observables: radius, height of the cone. Volume of cone corresponds to the reconstructed energy (E_{rec}).


Correlation between obtained fit parameters. LAQGSM

Initially we have experimental energy deposition E_{dep} in FHCaI.






After linear fit we have:

- E_{rec} is reconstructed energy (volume of cone);
- E_{max} – maximum energy in central bin (in FHCaI hole);
- Radius of spectator spot at FHCaI is defined by the scattering spot of spectators.




This correlation can be used for the centrality determination





Centrality resolution for E_{dep} vs E_{max}



Dependence of resolution of impact parameter on centrality




Centrality resolution for E_{dep} vs E_{max} 2% binning






Further approaches

2D linear fit method with subtraction of pion contribution



- **Narrow cone radius indicates that the outer FHCAL modules detect the pions mainly, while the spectators are detected by inner modules.**
- **Energy in outer modules can be regarded as pure non-spectator (pion) contribution.**
- **How to evaluate pion contribution in full FHCAL?**

Evaluation of pion energy contribution



- Linear fit with $y=kx+b$ background,
- b is known from outer FHCAL modules,
- k is taken from simulation and quite similar for LAQGSM and DCM-SMM models
- The ratio of edge and central energies is almost the same for different models (2.4609 for LAQGSM, 2.45876 for DCM-SMM)

Conclusion





- The ability of FHCAL to measure the collision centrality was considered.
- Only the spectators for the centrality reconstruction were used.
- Two + 1 methods for the centrality determination have been demonstrated:
 - Transverse-longitudinal energies correlation;
 - 2D-linear fit method;
 - Planned: 2D-linear fit with pion contribution subtraction method.
- A few new observables were introduced for the centrality determination.
- The usage of the introduced observables allows to determine the centrality more accurately, especially for the DCM-SMM model.
- DCM-SMM model provides worse centrality resolution because this model has much more heavy fragments which escape in FHCAL beam hole.
- The subtraction of the pion contribution probably makes possible to measure the energy of free (protons/neutrons) spectators. Number of free spectators can be estimated more accurately. It can be used for the centrality measurements.

Thank you for your attention!


This work was supported by the RFBR 18-02-40065 mega grant

BACKUPS

Energy deposition can be decomposed in two components: energy of free spectators and non-spectators energy




By using the subtraction of the non-spectator's contribution, the energy deposition can be decomposed into two components.





Both energies can be used for centrality determination.

E_pions vs Imp




Comparison LAQGSM 11 GeV







LAQGSM 11 GeV

Energy in the central bin vs impact parameter



MINUS BACKGROUND





Spectators scattering angle vs impact parameter



After subtracting the pion contribution, the energies for the central events become less

Centrality resolution for E_{dep} vs E_{max}

(after subtraction of pion contribution) backup







Dependence of resolution of impact parameter on centrality







Dependence of impact parameter on centrality


Comparison DCM-SMM 11 GeV бэкап




5 GeV example for LAQGSM and DCM-SMM models



LAQGSM and DCM-SMM models comparison for 5 GeV Erec Edep




Dependence of impact parameter on centrality





Dependence of resolution of impact
parameter on centrality



2D fit method LAQGSM 11 GeV



Correlation between transverse and longitudinal energies in FHCaI DCM-SMM 11 GeV backup



The separation of central and peripheral events with this model is clearly worse. This approach is not suited for DCM-SMM model



New approaches are needed

2d fit method results LAQGSM 11 GeV backup

