The 5th international conference on particle physics and astrophysics



Contribution ID : 903

Type : Oral talk

Nuclear modification factor in Pb+Pb collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV using Boltzmann's transport Equation with Tsallis statistics

Thursday, 8 October 2020 11:15 (15)

In this work the transverse momentum spectra and nuclear modification factor for heavy ion collisions are derived using Tsallis non-extensive statistics in relaxation time approximation. The Boltzmann-Gibbs Blast Wave (BGBW) function is used as the equilibrium function and Tsallis function is used as the initial distribution function while solving the Boltzmann transport equation in the relaxation time approximation. The framework is used to analyse the experimental data for particles like pions, kaons, protons, D^0 meson and J/ψ produced in Pb+Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV at the LHC, CERN. We find that our proposed equation of state describes the experimental data successfully.

Primary author(s): PAREEK, Pooja (Variable Energy Cyclotron Centre (IN)); TIWARI, Swatantra Kumar

Presenter(s) : PAREEK, Pooja (Variable Energy Cyclotron Centre (IN)); TIWARI, Swatantra Kumar **Session Classification :** Heavy Ion Physics

Track Classification : Heavy Ion physics