



Contribution ID : 671

Type : Oral talk

## Feasibility of thermal photon measurements in heavy ion collisions at NICA energies

*Tuesday, 6 October 2020 17:50 (15)*

Thermal photons serve as valuable probes of the hot and dense medium produced in heavy ion collisions. The effective thermal photon temperature measured at RHIC and LHC energies far exceeds the temperature predicted for the phase space transition into the deconfined state of quarks and gluons, known as quark-gluon plasma (QGP). Direct photon measurements in heavy ion collisions at the future NICA collider may help to measure the effective temperature of the produced medium at lower energies and trace the transition from QGP to the hadron gas state. In this contribution, we present feasibility studies on the thermal photon measurements in Au-Au collisions using the photon conversion method in the MPD experiment at NICA.

This work was funded by RFBR according to the research project No18-02-40045 and partially supported by the National Research Nuclear University MEPhI in the framework of the Russian Academic Excellence Project (contract No.02.a03.21.0005, 27.08.2013).

**Primary author(s) :** Dr. IVANISHCHEV, Dmitry (NRC «Kurchatov Institute» - PNPI); Dr. KOTOV, Dmitry (NRC «Kurchatov Institute» - PNPI); Dr. KRYSHEN, Evgeny (NRC «Kurchatov Institute» - PNPI); Dr. MALAEV, Mikhail (NRC «Kurchatov Institute» - PNPI); Dr. RIABOV, Victor (NRC «Kurchatov Institute» - PNPI); Dr. RYABOV, Yuriy (NRC «Kurchatov Institute» - PNPI)

**Presenter(s) :** Dr. IVANISHCHEV, Dmitry (NRC «Kurchatov Institute» - PNPI)

**Session Classification :** Heavy Ion Physics

**Track Classification :** Heavy Ion physics