The 5th international conference on particle physics and astrophysics



Contribution ID: 739

Type : Oral talk

Measurement of γ (ϕ_3) at Belle II

Wednesday, 7 October 2020 11:40 (15)

The CKM angle γ is the least well known of the angles of the unitarity triangle and the only one that is accessible with tree-level decays in a theoretically clean way. The key method to measure γ is through the interference between $B^+ \to D^0 K^+$ and $B^+ \to \overline{D}^0 K^+$ decays which occurs if the final state of the charmmeson decay is accessible to both the D^0 and \overline{D}^0 mesons. The Belle II experiment at the SuperKEKB energyasymmetric e^+e^- collider is a substantial upgrade of the B factory facility at the Japanese KEK laboratory. The design luminosity of the machine is $8 \times 10^{35} \text{ cm}^{-2} \text{s}^{-1}$ and the Belle II experiment aims to record 50 ab^{-1} of data, a factor of 50 more than its predecessor. Main operation of SuperKEKB has started in March 2019 and a results from the full available Belle II data set will be presented. To achieve the best sensitivity, a large variety of D and B decay modes is required, which is possible at Belle II experiment as almost any final state can be reconstructed including those with photons. With the ultimate Belle II data sample of 50 ab⁻¹, a determination of γ with a precision of 1 degree or better is foreseen. This talk will explain the details of the planned measurement at Belle II and include results related to these measurements obtained with the data already collected, including the first studies of the golden mode for ϕ_3 at Belle II: $B^+ \to D(K_S^0 \pi^+ \pi^-)K^+$.

Primary author(s) :PERUZZI, Ida Marena (INFN, LNF)Presenter(s) :PERUZZI, Ida Marena (INFN, LNF)Session Classification :High Energy Physics

Track Classification : High energy physics