Contribution ID : 178

Modeling of two-particle femtoscopic correlations at top RHIC energy

Tuesday, 11 October 2016 14:10 (20)

The spatial and temporal characteristics of particle emission source in high-energy collisions can be measured by using two-particle femtoscopic correlations. These correlations arise due to quantum statistics, Coulomb and strong final state interactions. In this talk, we report on the measurement of like-sign meson femtoscopic correlations produced in p+p, d+Au, Au+Au at top RHIC energy using Quantum Molecular Dynamics Model (UrQMD). Three-dimensional correlation functions are constructed using the Bertsch-Pratt parametrization of the two-particle relative momentum. The correlation functions are studied in several transverse momentum ranges. The emitting source radii of charged pions and kaons, R_{out} , R_{side} , R_{long} , are obtained from Gaussian fit to the correlation functions and compared to data from the STAR experiment.

Primary author(s) : Mr. NIGMATKULOV, Grigory (National Research Nuclear University MEPhI (Moscow Engineering Physics Institute)); Mr. ERMAKOV, Nikita (National Research Nuclear University MEPhI (Moscow Engineering Physics Institute))

Presenter(s) : Mr. ERMAKOV, Nikita (National Research Nuclear University MEPhI (Moscow Engineering Physics Institute))

Session Classification : Heavy Ion Student Session-1

Track Classification : Nuclear physics and particle physics