Contribution ID : 50

## The comparison of calculated atmospheric neutrino spectra with measurement data of IceCube and ANTARES experiments

Wednesday, 12 October 2016 15:30 (30)

The processing of the IceCube experiment data obtained during 988 days (2010–2014) revealed 54 neutrinoinduced events with deposited energies 20 TeV - 2 PeV [1]. The hypothesis of an astrophysical origin of these neutrinos is confirmed at 5.7 $\sigma$  CL. To identify reliably the neutrino events a thorough calculation of the atmospheric neutrino background should be performed. We calculate the atmospheric neutrino spectra in the energy range of 100 GeV - 10 PeV using the set of the hadronic models and several parametrizations of cosmic ray spectra supported by experimental data. It is shown that rare decays of short-lived neutral kaons contribute close to one third of the atmospheric conventional electron neutrinos at the energies above 100 TeV. The account for kaons production in pion-nucleus collisions gives rise to increase the  $\nu_e$  flux by 5–7% in the energy range of 100 GeV – 100 TeV. The detailed comparison of our calculations performed with use of Z(E, h) -functions approach [2], with those of MCEq method by A.Fedynitch et al. [3], shows the consistency on the whole at least in the energy range 100 GeV – 1 PeV. Calculated neutrino spectra agree rather well with the measurement data of the experiments IceCube [4,5] and ANTARES [6]. Uncertainties of the measurement data above 400 TeV leave a window for the the QGSM prompt neutrino component [2].

[1] Aartsen M.G. et al. (IceCube Collaboration). Evidence for high-energy extraterrestrial neutrinos at the IceCube detector // Science 2013. V. 342, 1242856; Phys. Rev. Lett. 113, 101101 (2014); arXiv:1510.05223.

[2] Sinegovskaya T.S, Morozova A.D., Sinegovsky S.I. High-energy neutrinos fluxes and flavor ratio in the Earth's atmosphere. Phys. Rev. D. 2015. V. 91, 063011.

[3] Fedynitch A. et al. Calculation of conventional and prompt lepton fluxes at very high energy, EPJ Web Conf. 2015. V. 99, 08001; arXiv:1503.00544.; Fedynitch A. et al. MCEQ - numerical code for inclusive lepton flux calculations. PoS (ICRC2015) 1129; http://github.com/afedynitch/MCEq/.

[4] Aartsen M.G. et al. (IceCube Collaboration). Development of a general analysis and unfolding scheme and its application to measure the energy spectrum of atmospheric neutrinos with IceCube. Eur. Phys. J. C. 2015. V. 75, 116.

[5] Aartsen M.G. et al. (IceCube Collaboration). Measurement of the atmospheric ve spectrum with IceCube. Phys. Rev. D. 2015. V. 91, 122004.

[6] Adrian-Martinez S. et al. Measurement of the atmospheric muon neutrino energy spectrum from 100 GeV to 200 TeV with the ANTARES telescope. Eur. Phys. J. C. 2013. V.73, 2606.

**Primary author(s) :** Ms. MOROZOVA, Anna (Moscow State U.); Prof. SINEGOVSKY, Sergei (Irkutsk State University, Institute of Apliied Physics)

**Co-author(s) :** Dr. KOCHANOV, Alexey (ISTP,SB RAS); Dr. SINEGOVSKAYA, Tatyana (Irkustk State Rail Way University)

**Presenter(s):** Ms. MOROZOVA, Anna (Moscow State U.); Prof. SINEGOVSKY, Sergei (Irkutsk State University, Institute of Apliied Physics)

Session Classification : Poster session - III

Track Classification : Nuclear physics and particle physics