

Study of LHC experiments sensitivity to anomalous quartic gauge couplings in $Z\gamma\gamma$ production during Run2

Monday, 10 October 2016 15:15 (30)

Run2 sensitivity to quartic anomalous gauge couplings was estimated for ATLAS experiment at LHC with increased energy of proton-proton collisions $\sqrt{s} = 13$ TeV and expected 40 fb^{-1} (2016) and 100 fb^{-1} (2016-2017) of integral luminosity. Simulation of $Z\gamma\gamma$ process with anomalous $ZZ\gamma\gamma$ and $Z\gamma\gamma\gamma$ couplings was performed using VBFNLO MC generator. Differential distributions on four body invariant mass of final state particles was used for extraction of expected limits on Effective Field Theory parameters f_{T0}/Λ^4 , f_{T5}/Λ^4 , f_{T9}/Λ^4 , f_{M2}/Λ^4 , f_{M3}/Λ^4 . Combined limits are obtained from two charged leptonic decay channels of Z boson ($Z\gamma\gamma \rightarrow l^+l^-\gamma\gamma$, where $l = e$ or μ). Unitarity of expected limits was studied using dipole form factor.

Primary author(s) : KUROVA, Anastasia (NRNU MEPhI); SOLDATOV, Evgeny (MEPhI)

Presenter(s) : KUROVA, Anastasia (NRNU MEPhI)

Session Classification : Poster session - I

Track Classification : Nuclear physics and particle physics