



Contribution ID : 317

Type : Oral talk

Low-energy spectra of nobelium isotopes: Skyrme random-phase-approximation analysis

Thursday, 24 October 2024 10:45 (15)

The low-energy multipole spectrum in isotopes 250-260No is investigated in the framework of fully self-consistent Quasiparticle-Random-Phase-Approximation (QRPA) method with Skyrme forces [1,2]. The representative set of Skyrme parametrizations (SLy5, SLy6, SkM* and SVbas) is applied. The main attention is paid to nuclei 252No and 254No, where we have most of the experimental spectroscopic information [3,4]. In addition to low-energy one-phonon collective states ($l\pi=20,22,30,31,32$) and their rotational band, the isomeric states are inspected. In general, a good agreement with the experimental data is obtained. Some K-isomers in these nuclei are inspected. It is shown that, in the chain 250–260No, features of 252No and 254No exhibit essential irregularities caused by a shell gap in the neutron single-particle spectra and corresponding break of the neutron pairing. The low-energy pairing-vibrational $K\pi = 0^+$ state is predicted.

[1] P.-G. Reinhard, B. Schuetrumpf, and J. A. Maruhn, *Comp. Phys. Commun.* 258, 107603 (2021).

[2] A. Repko, J. Kvasil, V.O. Nesterenko and P.-G. Reinhard, arXiv:1510.01248[nucl-th].

[3] R.-D. Herzberg and P.T. Greenlees, *Prog. Part. Nucl. Phys.* 61, 674 (2008).

[4] R.-D. Herzberg, arXiv:2309.10468[nucl-ex].

[5] F.L. Bello Garrote et al, *Phys. Lett. B*834, 137479 (2022).

Primary author(s) : MARDYBAN, Mariia

Presenter(s) : MARDYBAN, Mariia

Session Classification : Nuclear

Track Classification : Nuclear physics