The 7th international conference on particle physics and astrophysics



Contribution ID : 251

Type : Poster

Comparison of the 213Po half-life results measured by double and triple coincidences methods

Thursday, 24 October 2024 16:30 (30)

Method of measurement and result of processing of the 213Po α -active isotope half-life data measured in longrun continuous measurement with the underground low-background TAU-3 set-up are described. The set-up consists of two scintillation NaI(Tl) 150x150 mm detectors and double-layer (hlayer=1 mm) plastic scintillator detector (PCD) with d=18 mm. Source of the ...213Bi→213Po... decays placed between the PSD layers. The half-life calculated from a decay curve. The curve constructed from delay values between β - and α -pulses detected by the PSD. Two methods used for the event selection. The PSD pulses coincided in 16 mcs time window selected in the first case (double coincidences). Additional pulse of the NaI-SD detected γ - quantum from the 213Bi decay used in the second case for a validation of the 213Po birth and decay (triple coincidences). The values T1/2=3.6970±0.0005 mcs for the double coincidences and T1/2=3.6772±0.0005 (statistic)±0.005 (system) mcs for the triple coincidences were obtained 213Po half-life. Possible reasons of the result difference were discussed.

Primary author(s): KUZMINOV, Valery (INR RAS); KAZALOV, Vladimir (BNO INR RAS)
Presenter(s): KUZMINOV, Valery (INR RAS); KAZALOV, Vladimir (BNO INR RAS)
Session Classification: Poster session

Track Classification : Nuclear physics