



Contribution ID : 208

Type : Oral talk

## Double-hit experimental approach in studies of the multibody decays of heavy nuclei.

*Wednesday, 23 October 2024 10:45 (15)*

In our previous publications [1–3], a very specific effect, unknown in the past, was discussed, namely a break-up of the fission fragment while it passes through a solid-state foil. The fraction of the fragments which undergo the break-up is supposed to be born in the shape isomer states. The bulk of the results were obtained in the frame of the so-called missing mass experimental method when only one of two partners of the break-up is detected by the spectrometer. The difference between the total mass of the detected fragments and the mass of the mother system serves a sign of at least ternary decay. Alternative, so called” double-hit” approach lets obtain more direct information about the process. By definition, the double-hit registration approach means that two fragments were detected in the same PIN diode during one registration gate of 200 ns length. If a minimum time interval between their time stamps is less than 30 ns a pile-up of the signals take place. Restoring original signals from pile-up is discussed.

**Primary author(s) :** Mr. KAMANIN, D.V. (JINR); Mr. PYATKOV, Yu.V. (JINR,MEPHI); Mr. ZHUCHKO, V.E. (JINR); Mrs. GORYAINOVA, Z.I. (JINR); Ms. KUZNETSOVA, E.A. (JINR); Mr. SEREDA, Yu. M. (JINR); Mr. SOLODOV, A.N. (JINR); Mr. STREKALOVSKY, O.V. (JINR); Ms. ZHUKOVA, A.O. (JINR)

**Presenter(s) :** Mrs. GORYAINOVA, Z.I. (JINR)

**Session Classification :** Nuclear

**Track Classification :** Nuclear physics