

Contribution ID : 123 Type : Oral talk

Small- p_T production of J/ψ mesons within Soft Gluon Resummation Approach

Friday, 25 October 2024 10:15 (15)

Prompt heavy quarkonium production is well described within the collinear parton model in the next-to-leading order of perturbaive QCD at kinematical region of $p_T\gg M$ where M is a mass of a quarkonium state. But the region of small p_T is still being researched and the factorisation approach which is valid here is TMD-factorisation (transverse-momentum dependent). We studied J/ψ production in collision of protons at $\sqrt{s}=200,27$ and 19.4 GeV. Certainly, we used Nonrelativistic QCD (NRQCD) as a standard hadronisation model for charmonium production, and we extracted nonperturbative matrix elements for octet color states of the NRQCD from a set of experimental data on prompt J/ψ production because color singlet model can't be considered as sufficient for experimental data description.

The TMD approach is a general factorisation model for $p_T \ll M$ region [1]. One of the realisations of the TMD-factorisation is a so called Soft Gluon Resummation approach [2] where soft gluon emission by partons is considered, evolution of the TMD parton distribution functions is controlled by the Collins – Soper equations [3] though the TMD distribution is partly reduced to the collinear one. We perform our study at the LO+LL approximation for now. We describe data from PHENIX Collaboration at $\sqrt{s}=200$ GeV and from NA3 Collaboration at $\sqrt{s}=19.4$ GeV and do predictions for future SPD NICA experiments at $\sqrt{s}=27$ GeV. We considered both gluons and quarks as initial partons, we also estimated a contribution of P-wave charmonium production and calculated polarised J/ψ production as an angular coefficient λ . The Inverse-Error Weighting Scheme [4] is used as an approach for matching of collinear and TMD factorisations.

References:

[1] J. C. Collins. Foundations of Perturbative QCD, Cambridge Univ. Press, Cambridge (2011).

[2] J. Bor and D. Boer. TMD evolution study of the $\cos 2\phi$ azimuthal asymmetry in unpolarized J/ψ production at EIC, Phys. Rev. D 106, 1 (2022).

[3] J. C. Collins, D. E. Soper, and G. Sterman. Transverse momentum distribution in Drell-Yan pair and W and Z boson production, Nuclear Physics B 250 (1-4), 199–224 (1985).

[4] M. G. Echevarria, T. Kasemets, J.-P. Lansberg, C. Pisano, A. Signori. Matching factorization theorems with an inverse-error weighting, Phys. Lett. B 781, 161–168 (2018).

Primary author(s): SHILYAEV, Kirill (Samara University); SALEEV, Vladimir (Samara National Research

University)

Presenter(s): SHILYAEV, Kirill (Samara University)

Session Classification: HEP Theory

Track Classification: High energy physics: theory