The 7th international conference on particle physics and astrophysics



Contribution ID : 88

Type : Oral talk

Associated J/ψ and photon production in the Parton Reggeization Approach at high energy

Friday, 25 October 2024 10:00 (15)

The study of the associated production of J/ψ -mesons and photons is very impotent to verify perturbative quantum chromodynamics, factorization approaches, heavy quark to heavy quarkonium hadronization models and to extracting the Parton Distribution Function (PDF) of gluon in a proton from experimental data. In this paper, we study the associated production of J/ψ -mesons and photons at the energies of the Large Hadron Collider using the Parton Reggeization Approach (PRA) [1], which is based on Lipatov's effective field theory [2] and the modified Kimber-Martin-Ryskin model for unintegrated PDFs [3, 4]. We use two different heavy-quark to heavy quarkonium hadronization models: the non-relativistic quantum chromodynamics (NRQCD) [5] and the improved color evaporation model (ICEM) [6].

We have found that the PRA based predictions for production cross sections are higher than next-to-leading order collinear parton model results. We have shown that contributions of the direct production processes via the color-octet intermediate states in the NRQCD are suppressed relatively to the process via the color-singlet intermediate state, so the Color Singlet Model (CSM) can be used for predictions at the $p_{TJ/\psi} < 40$ GeV. The long distance matrix elements of the NRQCD and ICEM parameter F_{ψ} in the calculations are taken as they have been obtained to describe experimental data of the J/ψ -meson prompt production at the LHC energies.

We have predicted various differential cross-sections for associated production of J/ψ -mesons and photons in the PRA at the center-of-mass energy $\sqrt{s} = 13$ TeV in the central rapidity region. We have obtained that the PRA using the CSM cross-sections strongly overestimate the results obtained in the PRA using the ICEM and this difference increases as more as the cutoff on the photon transverse momentum p_T becomes larger. The results of this study indicate that there are big difference between the ICEM based and the NRQCD based predictions for the associated $J/\psi + \gamma$ production that can be used for discrimination heavy quark to heavy quarkonium hadronization models.

[1] M.A. Nefedov, V.A. Saleev, and A.V. Shipilova. Dijet azimuthal decorrelations at the LHC in the parton Reggeization approach. Physical Review D, 87(9):094030, 2013.

[2] L.N. Lipatov. Gauge invariant effective action for high energy processes in QCD. Nuclear Physics B, 452(1-2):369–397, 1995.

[3] M.A. Kimber, A.D. Martin, and M.G. Ryskin. Unintegrated parton distributions. Physical Review D, 63(11):114027, 2001.

[4] M.A. Nefedov and V.A. Saleev. High-energy factorization for the Drell-Yan process in pp and $p\bar{p}$ collisions with new unintegrated PDFs. Physical Review D, 102(11):114018, 2020.

[5] G.T. Bodwin, E. Braaten, and G.P. Lepage. Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium. Physical Review D, 51(3):1125, 1995.

[6] Y.Q. Ma, R. Vogt. Quarkonium production in an improved color evaporation model. Physical Review D, 94(11), 114029, 2016.

Primary author(s) : ALIMOV, Lev (Samara University); Dr. SALEEV, Vladimir (Joint Institute for Nuclear Research); KARPISHKOV, Anton (Joint Institute for Nuclear Research)

Presenter(s) : ALIMOV, Lev (Samara University)

Session Classification : HEP Theory

Track Classification : High energy physics: theory