

Accurate antineutrino spectra from fission fragments of ^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu

and some consequences followed from the spectra obtaining
procedure



V.V. Sinev, S.V. Ingerman

Institute for Nuclear Research RAS


Content

- Some history on antineutrino spectrum measurement (Rovno)
- High statistics spectrum measurements Double Chooz, Daya Bay, RENO
- What individual antineutrino spectrum should be used?
- Combined method of antineutrino spectrum calculation
- Spectra ^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu
- What cross section for ^{235}U should be?
- How accurate spectra could be used?


Rovno NPP experiment in 1988-90



Spectrometer used in Rovno on the assembling stage



V.I. Kopeikin and A.N. Kheruvimov



Spectrometer prototype ND-1

Spectrum and cross section by RONS and WIND

Statistics 174 000 neutrino events

$$\frac{dN_\nu}{dE} = 5.09 e^{-0.648E - 0.0273E^2 - 1.411(\frac{E}{8})^{10}}$$

^{235}U	^{238}U	^{239}Pu	^{241}Pu
0.586	0.075	0.292	0.047

Cross section WIND: $5.828 \pm 2.0\%$. Cross section RONS: $6.0 \pm 3.0\%$

Cross section WIND: $5.85 \pm 1.8\%$ (А.А. Кувшинников, Л.А. Микаэлян, С.В. Николаев, М.Д. Скорохватов, А.В. Этенко, Письма в ЖЭТФ, 54(5), 259-262, 1991)


^{235}U	^{238}U	^{239}Pu	^{241}Pu
0.614	0.074	0.274	0.038

Modern high statistics reactor neutrino experiments


Double Chooz



Experiment Double Chooz has been working in 2010-2017 in France (Ardenne). Red points mark the detector positions.





Detector Double Chooz scheme



Acrylic cylinders of the target and gamma-catcher of Double Chooz detector

Double Chooz spectrum measurement by near detector


~210 000 neutrino events



BKG smaller than for RONS and energy resolution is better

Measured spectrum ratio to the expected one calculated on base of Huber-Moeller.


Bump at ~ 5 MeV of visible energy exists in all experiments: Double Chooz, Daya Bay, RENO, NEOS et al.




How one can correctly calculate an antineutrino spectrum?

The standard is summation method.


The question – if we need to know exactly the individual fragment spectrum?



If the number of individual spectra is large – the shape of separate spectrum does not important





Which individual spectrum one should use?



The energy conservation law acts in the moment of particles production.

But later each particle has its individual fate.

Beta-decay



Distance at which W decays on electron and antineutrino, is about 10^{-14} cm . Neutron radius is estimated as $1.2-1.3 \cdot 10^{-13} \text{ cm}$.

This means that decay happens inside a neutron.

Usually a calculation uses data on decay-schemes keeping in databases. These data contain multiple branches of beta-decay.

How accurate these data?


Fission fragment yields are known with uncertainty estimated as
5%

We surely know branching for about 1/3 of all known fragments.
These are mainly long-lived nuclei.

For about 1/3 of fragments branching is totally unknown. These
nuclei are far from the line of beta-stability.

And about 1/3 of fragments have branching calculated, we can
say estimated.

Experimental antineutrino spectra weighted with IBD cross section




χ^2 square for minimization

$$\begin{aligned}\chi_k^2 = & \sum_i DC \frac{(y_{exp,i} - y_{calc,i})^2}{\sigma_i^2} + \sum_j DB \frac{(y_{exp,j} - y_{calc,j})^2}{\sigma_j^2} + \sum_l PROSPECT \frac{(y_{exp,l} - y_{calc,l})^2}{\sigma_l^2} + \\ & + \frac{(\sigma_{DC} - \sigma_{calc DC})^2}{\Delta\sigma_{DC}^2} + \frac{(\sigma_{DB} - \sigma_{calc DB})^2}{\Delta\sigma_{DB}^2} + \frac{(\sigma_{RENO} - \sigma_{calc RENO})^2}{\Delta\sigma_{RENO}^2} + \frac{(\sigma_{Bugey} - \sigma_{calc Bugey})^2}{\Delta\sigma_{Bugey}^2} + \\ & \frac{(\sigma_{Rovno} - \sigma_{calc Rovno})^2}{\Delta\sigma_{Rovno}^2}\end{aligned}$$


$$|\chi_k^2 - \chi_{k-1}^2| < \varepsilon$$

As a result of minimization we conform experimental antineutrino spectrum and obtain four spectra of ^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu . Obtained spectra perfectly describe all experimental spectra. Calculated on their base cross sections for core content of Double Chooz, RENO, Daya Bay, Bugey-3 have the smallest differences.

Antineutrino Spectra ^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu



What is experimental cross section of IBD?



$$\sigma_f = \int \frac{dn}{dE}(E) \sigma(E) dE$$

IBD cross sections for U and Pu isotopes in units 10^{-43} cm^2

	^{235}U	^{238}U	^{239}Pu	^{241}Pu
This work	5.917	10.95	4.220	6.347
ILL	6.426	8.929	4.204	5.796
Vogel	6.502	9.109	4.526	6.515
MEPhI	6.395	9.213	4.388	6.478
Huber & Mueller	6.681	10.12	4.387	6.081
Kopeikin et al.	6.308	9.395	4.33*	6.01*

before

$$\sigma_5 = 6.2 - 6.5$$

$$\sigma_8 = 9.58 - 10.50$$

$$\sigma_9 = 4.2 - 4.5$$

$$\sigma_1 = 5.95 - 6.1$$

now

$$\sigma_5 = 5.92$$


$$\sigma_8 = 10.95$$

$$\sigma_9 = 4.22$$

$$\sigma_1 = 6.35$$

What value of cross section for the spectrum of ^{235}U should be in reality ?

Antineutrino Spectra ^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu and mixed spectrum for Double Chooz



Mixed spectrum
practically coincides
with ^{235}U and ^{241}Pu

Measured with the best accuracy cross sections

	Experimental cross section
Double Chooz	5.74 ± 0.06
Daya Bay	5.84 ± 0.08
Bugey-4	5.752 ± 0.081
RENO	5.852 ± 0.094
WIND	5.85 ± 0.105

Fitting of Bugey-4 cross section

$$\sigma_f = \mathbf{5.752} \pm 0.081 = 0.538 \sigma_5 + 0.078 \sigma_8 + 0.328 \sigma_9 + 0.056 \sigma_9$$

$$\chi^2 = \frac{(\sigma - \sigma_{calc})^2}{\Delta\sigma^2}$$

$$\sigma_5 = \mathbf{5.91} \pm 0.77$$

$$\sigma_8 = 9.58 \pm 1.90$$

$$\sigma_9 = 4.52 \pm 0.59$$

$$\sigma_1 = 6.11 \pm 0.76$$

Results of fitting measured with best accuracy cross sections

Cross section	experimental	^{235}U	^{238}U	^{239}Pu	^{241}Pu
Double Chooz	5.74 ± 0.06 5.89	5.89 ± 0.63 5.97 ± 0.52	9.46 ± 1.73 10.0 ± 1.92	4.49 ± 0.75 4.60 ± 0.70	5.95 ± 0.62 6.38 ± 0.53
Daya Bay	5.84 ± 0.08	5.94 ± 0.56	9.85 ± 1.92	4.57 ± 0.72	6.36 ± 0.52
Bugey-4	5.752 ± 0.081	5.91 ± 0.77	9.58 ± 1.90	4.52 ± 0.59	6.11 ± 0.76
RENO	5.852 ± 0.094	5.94 ± 0.67	9.92 ± 1.63	4.58 ± 0.78	6.39 ± 0.51
WIND	5.85 ± 0.105	5.92 ± 0.78	9.72 ± 1.71	4.55 ± 0.59	6.49 ± 0.90
INR		5.92	10.95	4.22	6.35

Comparison of measured and calculated through INR spectra cross sections

experiment	Core content				$i\sigma_f \times 10^{-43}$	$^{INR}\sigma_f \times 10^{-43}$	R_{INR}	H-M	R_{H_M}
	^{235}U	^{238}U	^{239}Pu	^{241}Pu	[cm ² /fiss.]	[cm ² /fiss.]			
Double Chooz	0.520	0.087	0.333	0.060	5.71 ± 0.06	5.817	0.982	6.180	0.924
					5.74 ± 0.06		0.987		
Bugey-4	0.538	0.078	0.328	0.056	5.752 ± 0.081	5.779	0.995	6.163	0.933
Daya Bay	0.561	0.076	0.307	0.056	5.84 ± 0.08	5.804	1.006	6.204	0.927
RENO	0.571	0.073	0.300	0.056	5.852 ± 0.094	5.801	1.009	6.210	0.926

Conclusion


We applied new method of obtaining fission isotopes antineutrino spectra from high statistics experimental data.

Found four antineutrino spectra (^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu) allow to predict reactor antineutrino spectrum at any moment of its operation and at any core content.

Having exact (^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu) antineutrino spectra makes it possible to make online monitoring of reactor core during its cycle.


Thank you for the attention!

Деление тяжелых ядер происходит на два осколка неравной массы




Cr 60 24 0.49 s 0 0 2.6138e-12 1.0211e-11 3 1 2
 5701 0.102
 6059 0.886
 6111 0.012
 Mn 60 25 1.77 s 0 0 2.7318e-12 5.2985e-12 8 0.885 1
 5217 0.00503
 5229 0.0302
 5362 0.00591
 5522 0.02391
 5643 0.03523
 5923 0.81792
 6416 0.05537
 6601 0.02643
 Mn 60 25 0.28 s 0 0 3.0759e-13 5.9658e-13 4 1 1
 6088 0.02994
 6470 0.0499
 7621 0.04192
 8445 0.87824
 Fe 60 26 2.62e+06 y 0 0 0 0 1 1 1
 178 1
 Co 60 27 10.467 m 0 0 0 0 2 0.0025 1
 722.78 0.03459
 1548.88 0.95641
 Co 60 27 1925.28 d 0 0 0 0 2 1 1
 317.88 0.9988
 1492 0.12
 Ni 60 28 1e+30 y 0 0 0 0 0 1

The example of charge chain with mass A = 60



Анализ показал, что в базах данных только треть всех осколков имеет полностью известные схемы распада. Одна треть является полностью неизвестной, и еще одна треть имеет схемы распада, основанные на расчете по известным моделям ядра.

Массовое распределение осколков деления



Предлагаемый метод расчета:

Берем экспериментальный спектр или несколько

Рассчитываем спектр по базе данных

Вычисляем χ^2 . Если

Меняем схему распада у неизвестного ядра

Опять вычисляем χ^2 . Если χ^2 меньше предыдущего, то сохраняем нашу схему, если нет – восстанавливаем старую.


Переходим к следующему ядру.

Выполняем итерации пока разница χ^2 не станет меньше наперед заданного значения, например 10^{-5} .


Описываются экспериментальные точки (~25-30 точек) и получается χ^2 с большим количеством параметров

Пример варьирования вероятностей распада для ^{144}Ba

#	<E _β > [keV]	I _β (abs) [%]	Daughter level [keV]	J ^π	E _β , max [keV]	logft	Transition type	Comments
1	720.25	0.4	1240.19 10		(1879)	6.1		
2	727.25	1.3	1223.86 8		(1895)	5.6		
3	775.26	0.7	1117.05 9		(2002)	6.0		
4	789.26	2.6	1085.82 7	1+	(2033)	5.4	allowed	
5	832.26	9.2	990.61 5	1+	(2128)	4.9	allowed	
6	938.26	19.2	756.19 4	1+	(2363)	4.8	allowed	
7	945.26	< 0.6	740.60 6		(2378)	> 6.3		
8	977.26	< 0.3	669.88 5		(2449)	> 6.7		
9	984.26	< 0.3	655.19 5		(2464)	> 6.7		
10	1007.26	41.0	603.37 3	1+	(2516)	4.6	allowed	
11	1032.26	< 0.02	549.98 9		(2569)	> 7.9	1 st unique	
12	1065.26	1.0	477.45 6		(2642)	6.3		
13	1066.26	2.3	475.47 4	(1-)	(2644)	5.9	1 st non-unique	
14	1066.26	1.3	474.57 3	(1-)	(2644)	6.2	1 st non-unique	
15	1088.26	< 0.2	401.23 4	(2-)	(2718)	> 8.3	1 st unique	
16	1108.26	< 1.2	382.94 4	(1-,2-)	(2736)	> 6.3		
17	1146.26	2.2	300.68 4	(0-)	(2818)	6.1	1 st non-unique	
18	1149.26	< 0.8	294.72 4	(2-,1-)	(2824)	> 6.5		
19	1186.26	> 5.5	215.167 6	(1-)	(2904)	< 5.7	1 st non-unique	
20	1191.26	< 3.7	172.830 7	(2-)	(2946)	> 7.3	1 st unique	
21	1199.26	< 1.6	185.681 6	(1-)	(2933)	> 6.3	1 st non-unique	
22	1223.26	< 3.2	103.855 6	(2-)	(3015)	> 7.4	1 st unique	



We change Q the array of $T_{\max}$ for beta-decays



ENDF Radioactive decay data /MF8.MT457/

by V.Zerkin, IAEA-NDS, 2020-2022, ver.2022-08-29

☒ Show all spectra. [Hide](#) low-intensity lines ($\leq 1\%$).

[Select](#)/[unselect](#) data for plotting. [Plot](#) data in %: ☒ using normalization: ☒

☒ #1.ENDF/B-VIII.0 ☒ #2.JEFF-3.3 ☒ #3.JENDL-5 ☒ #4.ENSDF-2024


#1. ENDF/B-VIII.0 Ba-144	#2. JEFF-3.3 Ba-144	#3. JENDL-5 Ba-144	#4. ENSDF-2024 Ba-144
<p>Nucleus: Ba-144 ZA=56144 Library: ENDF/B-VIII.0, MAT=1855 AUTH: Conv. from CGM EDATE: EVAL-AUG11 Half life: $11.5 \pm 0.2(s)$ AWR: 142.6866 Isomer number: LISO=0 Level number: LIS=0 Spin & Parity: 0+ Ebeta: $902.2703 \pm 47.1652 (keV)$ Egamma: $785.05 \pm 33.37 (keV)$ Ealpha: $0 \pm 0 (keV)$ Decay modes: 1 Radiation types:2</p>	<p>Nucleus: Ba-144 ZA=56144 Library: JEFF-3.3, MAT=1864 AUTH: G.AUDI, O.BERSILLON, J.BLACHOT + EDATE: EVAL-DEC03 Half life: $11.5 \pm 0.2(s)$ AWR: 142.687 Isomer number: LISO=0 Level number: LIS=0 Spin & Parity: 0+ Ebeta: $929.8 \pm 9.298 (keV)$ Egamma: $784.7 \pm 7.847 (keV)$ Ealpha: $0 \pm 0 (keV)$ Decay modes: 1 Radiation types:0</p>	<p>Nucleus: Ba-144 ZA=56144 Library: JENDL-5, MAT=1971 AUTH: Conversion from ENSDF EDATE: EVAL-NOV21 Half life: $11.5 \pm 0.2(s)$ AWR: 142.6866 Isomer number: LISO=0 Level number: LIS=0 Spin & Parity: 0+ Ebeta: $929.8 \pm 34.4609 (keV)$ Egamma: $784.7 \pm 11.1107 (keV)$ Ealpha: $0 \pm 0 (keV)$ Decay modes: 1 Radiation types:2</p>	<p>Nucleus: Ba-144 ZA=56144 Library: ENSDF-2024 Half life: $11.5 \pm 0.2(s)$ Spin & Parity: 0+ Library ; ENSDF 202410 by livechart20241016 LARA Nuclide ; Ba-144 Element ; Barium Z ; 56 Daughter(s) ; (B-) ; La-144 ; 100 Q- ; 3083 Possible parent(s) ; Jp ; 0+ Half-life (s) ; 11.5 ; 0.2 Half-life (s) ; 1.15E1 ; 2E-1 Decay constant (1/s) ; 6.03E-2 ; 1E-3 Specific activity (Bq/g) ; 2.523E20 ; 4.184E18 Reference ; ENSDF 200108 / Author(s) A.A. Sonzogni Emissions (128 lines) sorted by increasing energy Decay modes: 1 Radiation types: 2</p>
<p>#M1. Decay mode: RTYP=1 β^- Decay Q=$3066.408 \pm 19.357 keV$</p>	<p>#M1. Decay mode: RTYP=1 β^- Decay Q=$3121 \pm 0 keV$</p>	<p>#M1. Decay mode: RTYP=1 β^- Decay Q=$3119 \pm 56 keV$</p>	<p>#M1. Decay mode: RTYP=1 β^- Decay Q=$3083 keV$</p>
<p>#R1. Radiation type: STYP=0 γ AveDecayEne=$586.72 \pm 0 keV$ ☒ ContinuousSpectrum:307 ☐ plot</p>	-no-	<p>#R1. Radiation type: STYP=0 γ AveDecayEne=$784.7 \pm 11.111 keV$ ☒ DiscreteSpectrum: 88 lines ☐ plot ☒ ContinuousSpectrum:428 ☐ plot</p>	<p>#R1. Radiation type: STYP=0 γ AveDecayEne: ? ☒ DiscreteSpectrum: 100 lines ☐ plot</p>
<p>#R2. Radiation type: STYP=1 β^- AveDecayEne=$993.65 \pm 0 keV$ ☒ ContinuousSpectrum:307 ☐ plot</p>	-no-	<p>#R2. Radiation type: STYP=1 β^- AveDecayEne=$929.8 \pm 1.4837 keV$ ☒ DiscreteSpectrum: 22 end-points ☐ plot ☒ ContinuousSpectrum:349 ☐ plot</p>	-no-
-no-	-no-	-no-	<p>#R2. Radiation type: STYP=9 X-rays AveDecayEne: ? ☒ DiscreteSpectrum: 6 lines ☐ plot</p>

Legend


left value	right value	ratio of values	display
Value1	Value2	Value1==Value2	Value2

Еще пример ^{75}Zn

$^{75}_{30}\text{Zn}$ - estimated, статус 2



Измеренные спектры антинейтрино и расчетные



Отношение измеренных спектров антинейтрино к нашему спектру

